Successful atrial fibrillation ablation without pulmonary vein isolation utilizing focal impulse and rotor mapping in an atriopulmonary Fontan



Madhukar Kollengode, MD,*[†] Jehu Mathew, MD,* Elizabeth Yeung, MD,*[†] William H. Sauer, MD, FHRS,* Duy Thai Nguyen, MD, FHRS*

From the *Division of Cardiovascular Medicine, University of Colorado, Anschutz Medical Center, Aurora, Colorado, and [†]Division of Cardiology, Children's Hospital Colorado, Aurora, Colorado.

Introduction

Fontan palliation is designed to separate systemic and pulmonary circulations to ameliorate systemic hypoxemia and relieve ventricular volume overload in patients with a functional univentricular heart. Since its introduction in 1971, the surgical techniques have evolved considerably.¹ Early atriopulmonary (AP) versions of the Fontan incorporated the right atrium (RA) into the systemic venous circuit and are associated with considerable burden of early and late atrial dysrhythmias.¹⁻⁵ Although the majority of atrial arrhythmias are macroreentrant RA circuits, such as intra-atrial reentrant tachycardia (IART), the prevalence of late atrial fibrillation (AF) in Fontan patients is increasing.^{2,4–7} The mechanisms responsible for AF initiation and maintenance in this population and the role and techniques of catheter ablation for treatment of AF are not well defined.^{6,8-12} We present a case of successful AF ablation without pulmonary vein (PV) isolation by utilizing focal impulse and rotor mapping to identify and target mechanisms of AF maintenance in a patient with atrial arrhythmias and medication intolerance in the setting of an AP Fontan.

Case report

A 25-year-old woman with complex congenital heart disease presented with recurrent palpitations and exertional syncope. She had a history of tricuspid atresia, initially palliated with a

KEYWORDS Adult congenital heart disease; Atrial fibrillation; Catheter ablation; Congenital heart disease; Focal triggers; Fontan procedure; Mechanisms

(Heart Rhythm Case Reports 2018;4:241-246)

Drs Sauer and Nguyen receive significant research and consulting grants from Biosense Webster and CardioNXT and educational grants from St. Jude Medical, Boston Scientific, and Medtronic. Drs Nguyen and Sauer have non-public equity interests/stock options in CardioNXT. No specific grant from funding agencies in the public, commercial, or not-for-profit sectors was utilized for the preparation of this manuscript. There are no disclosures related to the topic of this manuscript. **Address reprint requests and correspondence:** Dr Duy Thai Nguyen, Cardiac Electrophysiology, University of Colorado, 12631 East 17th Ave, Campus Box B132, Aurora, CO 80045. E-mail address: Duy.t.nguyen@ucdenver.edu. left modified Blalock-Taussig shunt and a subsequent shunt ligation and AP Fontan. Her childhood and adolescence were relatively unremarkable with the exception of an embolic stroke at age 20 while taking oral contraception, from which she had a complete neurologic recovery and for which she is maintained on anticoagulation. Because of progressive cyanosis and arrhythmias, her adult cardiologist referred her to a children's hospital, where catheterization revealed significant collateralization without focal hemodynamic derangement. A Fontan conversion was recommended, but she declined and shortly afterward relocated and was referred to our program.

On the initial visit, the patient reported significant functional impairment due to fatigue and daily symptomatic palpitations, lasting minutes to hours. Examination was notable for cyanosis and clubbing, with resting O_2 saturation of 83%. Echocardiography demonstrated tricuspid atresia with an unobstructed AP Fontan, severe RA dilation, and normal systemic ventricular function. Baseline electrocardiography demonstrated sinus rhythm with a prolonged PR interval of 216 ms, evidence of RA enlargement, and left-axis deviation. Exercise testing was notable for desaturation to 72%, poor exercise capacity (peak VO₂ 17.2 mL/kg/min), and salvos of a regular tachyarrhythmia with ventricular rate of 166 bpm (Figure 1A). Event monitoring revealed symptomatic paroxysmal AF (Figure 1B). Catheterization revealed unobstructed Fontan flow with mean resting Fontan pressure of 5-6 mm Hg. A large venoatrial collateral from the superior vena cava to the left atrium was identified and the coil embolized, with improvement in systemic saturation to 89%. Medical management with beta-blocker was not tolerated. Given her overall normal hemodynamics and her aversion to Fontan conversion or antiarrhythmics, the decision was made to attempt catheter ablation. Preprocedural imaging was performed with cardiac magnetic resonance imaging to evaluate the atrial anatomy and identify pulmonary venous structures.

Description of the procedure

The patient was brought to the electrophysiology laboratory in sinus rhythm. Intracardiac echocardiography was used to

KEY TEACHING POINTS

- Atrial tachyarrhythmias are a common source or morbidity in patients who undergo classic atriopulmonary (AP) Fontan. The prevalence of atrial fibrillation (AF) is increasing in this population.
- Catheter ablation of AF in AP Fontan patients can be feasible, safe, and efficacious.
- Factors driving AF in Fontan patients are ill defined, and nonpulmonary vein triggers should be considered. Pulmonary vein isolation may not be necessary to maintain sinus rhythm in AP Fontan patients with nonpulmonary vein triggers for AF.
- Ablative strategies targeted toward mechanisms of AF maintenance, such as rotors and focal impulses, should be considered in AP Fontan patients with AF.
- Meticulous identification of triggers and judicious, focused ablative approaches applied by electrophysiologists skilled in catheter ablation and familiar with complex congenital heart disease should be considered an option to treat drug refractory atrial arrhythmias in Fontan patients. A sustained reduction in arrhythmia burden and improvement in quality of life can be achieved in select patients.

guide catheter placement, to create an anatomic geometry of the Fontan pathway to merge with her preprocedural cardiac magnetic resonance imaging, and to monitor hemodynamics and ablation lesions. Therapeutic anticoagulation was maintained during the case. Various catheters, including a decapolar catheter, PentaRay multielectrode catheter (Biosense Webster, Diamond Bar, CA), and a ThermoCool Smart-Touch ablation catheter (Biosense Webster), were positioned throughout the Fontan pathway (Figure 1D). A dense voltage map of the Fontan showed mostly normal voltage, except for a dense strip of scar within the inferolateral limb of the Fontan, contiguous with the inferior vena cava (Figure 1E).

Electrophysiological testing was performed once the voltage map was acquired. Triggers initiating an IART with a tachycardia cycle length of 280–310 ms were noted during isoproterenol infusion (Figure 1C). Activation mapping demonstrated a microreentrant circuit with early activation within the inferolateral Fontan, at the border zone between normal tissue and scar. A series of ablation lesions delivered within the border zone resulted in slowing of tachycardia before termination and transition into AF. Given the persistence of AF, substrate modification was then performed, extending the ablation lesions from the region of IART termination superiorly and inferiorly through the border zone to transect channels within scar.

Furthermore, in order to identify possible drivers of AF maintenance, the decision was made to map potential rotors and focal impulses. Using a focal impulse and rotor mapping multielectrode basket catheter and RhythmView mapping system (Topera Inc, Abbott EP, Abbott Park, IL), rotors and focal impulses were identified. A focal impulse and an adjacent fixed rotor were identified in the lateral aspect of the Fontan pathway near the previously identified border zone of the Fontan scar (Figure 2). The region containing the rotor and focal impulse was ablated to noncapture, and further ablation was performed to anchor this region inferiorly to the previously identified area of scar. AF selfterminated. Bidirectional block was then confirmed across ablation lines. Several other rotors noted to be within areas of normal voltage or along the atrioventricular node/His region were not targeted for ablation given the potential risks of complete heart block and proarrhythmia.

An additional macroreentrant cavo-tricuspid isthmus (CTI)–dependent atrial flutter (AFL) circuit with a tachycardia cycle length of 230 ms was induced with rapid atrial pacing and isoproterenol infusion. CTI ablation was performed with AFL termination, and bidirectional block was confirmed with differential pacing. Repeat pacing failed to induce sustained atrial arrhythmias. The patient has remained free of significant arrhythmias without medications for over 18 months, with marked improvement in her functional capacity and quality of life.

Discussion

Atrial arrhythmias are highly prevalent in patients with Fontan palliation for functional univentricular heart and are associated with significant morbidity and mortality.^{2,5} Arrhythmias may arise from obstructive hemodynamic perturbations, progressive atrial dilation and remodeling, or scar. There is a high failure rate of antiarrhythmic drugs used to treat atrial arrhythmias in these patients.¹³ Catheter ablation of atrial arrhythmias in patients with Fontan palliation, the majority of which are right-sided, has historically been associated with lower success than in other forms of congenital heart disease, with recurrence rates of 30%-60% reported within 1 year.⁵ Ablation of atrial arrhythmias is challenging in Fontan patients for myriad reasons, including distorted anatomy resulting in anatomic and electrical barriers to native atrial tissue, restricted catheter access, inability to deliver lesions of sufficient penetrance, and hemodynamic instability during protracted cases.

AF ablation in Fontan patients is uncommon. Whether the optimal ablative strategy is to target all possible triggers utilizing empiric lines for IART prophylaxis and PV isolation or to selectively identify and target clinically relevant triggers is unclear. We and others have demonstrated durable AF ablation in Fontan patients by solely targeting right-sided triggers.^{7,11,14} AF recurrence rates are higher in patients with non-PV triggers for AF,¹² and non-PV triggers for AF, including right-sided triggers, are more common in patients with a normal-sized left atrium,¹¹ such as many

Download English Version:

https://daneshyari.com/en/article/8660473

Download Persian Version:

https://daneshyari.com/article/8660473

Daneshyari.com