

Utility of entrainment pacing to clarify the circuit of macroreentrant tachycardia with dual early sites on activation maps

Koichi Nagashima, MD, PhD, Yasuo Okumura, MD, PhD, Ryuta Watanabe, MD, Masaru Arai, MD, Yuji Wakamatsu, MD, Ichiro Watanabe, MD, PhD

From the Division of Cardiology, Department of Medicine, Nihon University School of Medicine, Tokyo, Japan.

Introduction

Three-dimensional (3D) activation mapping is useful for identifying the substrate of macroreentrant tachycardias, such as atrial and ventricular tachycardias, that are caused by macroreentry.¹⁻³ To identify the reentrant circuit, the activation map should account for the entire tachycardia cycle length (TCL).^{1,3} In some cases, the tachycardia cycle is longer than the total activation time (TAT) of the target chamber. In patients with a severely diseased heart and in those having undergone cardiac surgery or an ablation procedure, there can be bystander regions adjacent to the reentrant circuit that activate late, almost simultaneously with the next tachycardia beat. This phenomenon masks the actual reentrant circuit. We illustrate the utility of identifying orthodromic capture during entrainment pacing to distinguish the critical circuit of the macroreentrant tachycardia from the bystander region in a case of atrial tachycardia (AT) and a case of ventricular tachycardia (VT).

Case report

A case of AT

The patient was a 65-year-old man who had undergone ablation for persistent atrial fibrillation and required radiofrequency (RF) ablation for AT. The previous ablation procedure included pulmonary vein isolation (PVI) and creation of a left atrial (LA) roof line. The electrocardiogram showed AT with a P-wave morphology that was positive/negative in the inferior limb leads and positive in V1. An activation map was obtained during AT (TCL: 205 ms) with a

KEYWORDS Reentry; Three-dimensional mapping; Total activation time; Tachycardia cycle length; Entrainment pacing; Orthodromic capture (Heart Rhythm Case Reports 2017; ■:1-4)

The study was supported by departmental resources only, and the authors have no conflict of interest to declare. **Address reprint requests and correspondence:** Dr Koichi Nagashima, Division of Cardiology, Department of Medicine, Nihon University School of Medicine, 30-1 Ohyauguchikamicho, Itabashi-ku, Tokyo 173-8610, Japan. E-mail address: cocakochan@gmail.com.

KEY TEACHING POINTS

- Three-dimensional (3D) activation mapping for macroreentrant tachycardia when the total activation time is greater than the tachycardia cycle length (TCL) is challenging because there can be bystander regions that activate late, simultaneously with the next tachycardia beat within the reentrant circuit.
- In such tachycardias, 3D mapping often shows 2 or more sites of earliest activation because the bystander region is detected as a region of early activation owing to the window of interest.
- From electrophysiological studies of an atrial macroreentrant tachycardia and a ventricular tachycardia, we propose a method in which mapping catheters are separately positioned at the 2 sites of earliest activation and entrainment pacing is performed from both sites.
- This allows us to identify the site at which the post-pacing interval is identical to the TCL as the “true” site of earliest activation on the circuit and the site at which orthodromic capture of entrainment pacing from the “true” site occurs as the bystander site with “pseudo” earliest activation.

CARTO3 mapping system (Biosense Webster, Diamond Bar, CA) and use of a 20-pole, 5-spline catheter (4 mm inter-electrode spacing; PentaRay NAV; Biosense Webster). The right atrial (RA) activation map accounted for only 53% (108 ms) of the TCL, and the site of earliest activation was a wide RA septal area. The LA activation map accounted for 93% (190 ms) of the TCL and revealed 2 sites of earliest activation: the roof of the LA and the posterior aspect of the left superior pulmonary vein (LSPV, [Figure 1A](#)). An ablation




Figure 1 Electrocardiograms and 3-dimensional activation maps obtained in the case of atrial tachycardia (AT). **A:** Left atrial (LA) activation map of the AT. **B:** Entrainment pacing from the LA roof (pacing site 1, panel A). **C:** Entrainment from the other site of earliest activation in the left superior pulmonary vein (LSPV) (pacing site 2, panel A). **D:** The accurate activation map with the reannotation of the electrograms according to the tachycardia cycle length (TCL) and the results of entrainment pacing. The reentrant circuit is shown in red, yellow, green, and blue, and the bystander area is shown in pink. LA roof-dependent macroreentrant AT (black arrow) with a bystander LSPV region through the gap of the prior pulmonary vein isolation line (gray arrow) was diagnosed. ABL = ablation catheter; AP = anterior-posterior; CS = coronary sinus; LAA = left atrial appendage; LAT = local activation time; LIPV = left inferior pulmonary vein; MA = mitral annulus; PA = posterior-anterior; PV = pulmonary vein; RIPV = right inferior pulmonary vein; RSPV = right superior pulmonary vein; TAT, total activation time.

catheter and PentaRay catheter were positioned at the LA roof (Figure 1A, pacing site 1) and LSPV (Figure 1A, pacing site 2), respectively, and entrainment pacing from the LA roof (pacing site 1) revealed that the post-pacing interval (PPI) (215 ms) was almost identical to the TCL of 205 ms, suggesting that the LA roof was included in the circuit (Figure 1B). Furthermore, the last pacing stimulus from the LA roof also captured the LSPV 1 cycle after LA electrogram activation. In contrast, entrainment pacing from the LSPV (pacing site 2) failed to capture the LA potentials despite capture of the pulmonary vein potentials, suggesting unidirectional exit block at the prior PVI line (Figure 1C). LA roof-dependent macroreentrant AT was diagnosed, and delivery of RF energy to the LA roof eliminated the AT and was followed by repeat LSPV isolation. We created an accurate activation map by reannotation of the electrograms according to the TCL and the results of entrainment pacing. Correctly annotated, the LA TAT was 438 ms and greater than the TCL of 205 ms (Figure 1D).

A case of VT

The patient was a 79-year-old man in whom VT developed after he had undergone coronary stenting for treatment of

extensive anteroseptal myocardial infarction and then endoventricular circular patch plasty (Dor procedure) for a left ventricular (LV) apical aneurysm. The QRS complex morphology was that of left bundle branch block with an undetermined axis and a TCL of 350 ms. Right ventricular (RV) and LV activation mapping showed a centrifugal activation pattern with 2 sites of earliest activation: the RV apical septum and the LV apical septum. TAT of both ventricles on the 3D map accounted for 45% (156 ms) of the TCL, suggesting a focal VT mechanism rather than macroreentry (Figure 2A). However, entrainment pacing from the RV apex (site of earliest activation) showed concealed entrainment and a PPI-TCL of 5 ms with a short stimulus-QRS interval of 15 ms (Figure 2B). Concealed entrainment with a PPI identical to the TCL and a long stimulus-QRS interval of 320 ms was also obtained by pacing from the LV apex (the other site of earliest activation) opposite the RV site (Figure 2C). Furthermore, the last pacing stimulus also captured the basal aspect of the LV (LV potentials in the coronary sinus), as well as the QRS, until the next tachycardia beat at the LV apex, indicating that LV apical potentials were activated 1 cycle before the QRS and the basal LV. With this phenomenon and the bipolar voltage map indicating an extensive low-voltage LV apical scar, we presumed

Download English Version:

<https://daneshyari.com/en/article/8660523>

Download Persian Version:

<https://daneshyari.com/article/8660523>

[Daneshyari.com](https://daneshyari.com)