

# A surgeon with symptomatic ventricular tachycardia while operating and exacerbated by left lateral decubitus position: Too much of a stretch?

Faris Khan, MD, MS,\* Gregory K. Feld, MD, FHRS,\* Amir Schricker, MD, FHRS,\*  
Frederick T. Han, MD, FHRS,<sup>†</sup> Kurt S. Hoffmayer, MD, FHRS,\* Jonathan C. Hsu, MD, MAS, FHRS\*

From the \*Section of Cardiac Electrophysiology, Division of Cardiology, University of California San Diego, La Jolla, California, and <sup>†</sup>Section of Cardiac Electrophysiology, Division of Cardiology, University of Utah, Salt Lake City, Utah.

## Introduction

Idiopathic ventricular arrhythmias are seen in patients with otherwise structurally normal hearts. Common areas of origin include the ventricular outflow tracts, the left ventricular (LV) base including the intracavitary structures, and the right ventricular (RV) inflow region. The moderator band (MB) of the RV is an intracavitary structure that has been implicated in idiopathic ventricular arrhythmias.<sup>1,2</sup> The authors report a case of an otherwise healthy surgeon with palpitations and light-headedness associated with premature ventricular contractions (PVCs) and ventricular tachycardia (VT), exacerbated by the left lateral decubitus position and a potential stretch mechanism, and found to be arising from the RV MB.

## Case report

A 62-year-old neurosurgeon with no significant medical history was in his usual state of health until a week prior when he began having intermittent palpitations. Subsequently, while standing and performing a surgical procedure in the operating room, he felt the sensation of palpitations and profound light-headedness. He was unable to complete the surgical procedure, and staff in the operating room were able to perform a 12-lead electrocardiogram (ECG; in the supine position), which showed frequent PVCs and 2-lead telemetry showed salvos of nonsustained ventricular tachycardia (NSVT) of the same morphology (Figures 1A and 1B). The patient denied any other symptoms of angina or acute coronary syndrome, including chest pain, shortness of breath, paroxysmal nocturnal dyspnea, or orthopnea. He

**KEYWORDS** Moderator band; Ventricular tachycardia; Ablation; Right ventricle; Stretch; Premature ventricular contractions (Heart Rhythm Case Reports 2017;0:1–5)

**Address reprint requests and correspondence:** Dr Jonathan C. Hsu, Section of Cardiac Electrophysiology, Division of Cardiology, University of California San Diego, 9454 Medical Center Dr 3rd Fl, Rm 3E-417 La Jolla, CA 92037. E-mail address: Jonathan.Hsu@ucsd.edu.

denied drug use and had no family history of cardiac disease or sudden cardiac death. Physical examination was normal. Laboratory examination was notable for normal potassium and creatinine levels.

After admission during hospitalization, he continued to have palpitations and profound light-headedness associated with PVCs and NSVT of up to 30 beats in length (Figure 1B), despite treatment with intravenous lidocaine. He underwent coronary angiography, which showed severe disease involving the first diagonal branch of the left anterior descending artery, but the remaining arteries did not have significant obstructive coronary artery disease (CAD) (Online Supplemental Video 1). Because of the sudden onset of PVCs and VT and no overt clinical symptoms of angina or acute coronary syndrome, percutaneous coronary intervention of the diagonal branch was deferred and medical management was advised by the interventional cardiology team for newly diagnosed CAD. His transthoracic echocardiogram showed normal LV and RV function with preserved ejection fraction and no LV wall motion abnormalities. Upon further questioning, he reported noticing distinct and reproducible worsening of ventricular ectopy including ventricular bigeminy and NSVT while on cardiac telemetry monitoring when he was placed in the left lateral decubitus position. Upon lying in the right lateral recumbent or supine position, he had minimal ventricular ectopy. Because of the recurrent nature of his ventricular arrhythmias, an electrophysiology study and possible ablation was recommended after a cardiac magnetic resonance imaging was performed. Cardiac magnetic resonance imaging showed focal hypokinesis and delayed enhancement in the mid- and apical anterior LV myocardium (diagonal artery territory) compatible with a prior ischemic event and no other abnormality (Supplemental Video 1). LV and RV function was overall preserved.

## Analysis of the ECG

The patient's baseline ECG showed sinus rhythm with PVCs in a bigeminal pattern (Figures 2A and 2B). A review of the

**KEY TEACHING POINTS**

- Stretch mechanism can be involved in ventricular tachycardia induction from the moderator band.
- Use of intracardiac echocardiography can be useful to confirm ablation catheter position around the moderator band insertion site.
- Idiopathic ventricular tachycardia originating from the moderator band may have excellent outcomes after radiofrequency ablation.

telemetry showed that all ventricular ectopy was monomorphic with a left bundle branch block morphology with superior axis, and the 12-lead ECG showed an R-wave transition from negative to positive at precordial lead V<sub>5</sub>, left bundle branch block pattern, and left superior axis. Vectorcardiography indicated an inferolateral RV free wall exit site.

**Electrophysiology study**

In the electrophysiology laboratory, the patient was prepped and draped in a standard fashion, and 3-dimensional electroanatomic mapping patches and 12-lead ECG patches were placed. While the patient was in the supine position, there was no evidence of ventricular arrhythmias, even without sedative medication. Subsequently, provocative testing was performed by placing the patient in the left lateral decubitus position, which immediately reproduced the clinically relevant ventricular ectopy in a bigeminal pattern (Figures 2A–2D). The ventricular ectopy was recorded, and a 12-lead ECG morphology template was created for possible pace-map matching with the recording system (Bard Electrophysiology, Lowell, MA). The CARTO 3-dimensional electroanatomic mapping system (Biosense Webster Inc., Diamond Bar, CA) was used with the CARTO-Sound module along with an 8-F phased array CARTO-Sound intracardiac echocardiography (ICE) catheter. CARTO-Sound and ICE were used to precisely delineate the anatomy of the RV endocardium, papillary muscles, and MB. A 3.5-mm D/F curve Smart TouchThermocool (Biosense Webster) mapping/ablation catheter was used. After creating geometry using the mapping catheter in addition to CARTO-Sound, we attempted to induce VT using burst pacing from the RV apex down to 300 ms and programmed electrical stimulation at 500 ms with single extrastimuli down to 300 ms. No ventricular ectopy or VT could be induced with RV pacing at baseline or during infusion of isoproterenol up to 5 µg/min. Atrial burst pacing was also performed from the proximal coronary sinus catheter down to a cycle length of 400 ms; however, this was also not able to induce ventricular arrhythmia. Owing to a lack of ventricular ectopy in the supine position despite ventricular pacing and programmed electrical stimulation, we used the previously templated PVCs obtained when the patient was in the left lateral decubitus position to perform pace mapping in an attempt to localize the exit site

of a presumed ectopic ventricular arrhythmia source. Under guidance of a 3-dimensional geometry created by CARTO-Sound, an RV MB was identified that extended from the inferior-lateral and mid free wall of the RV to the basal septum of the RV. Pace mapping was performed from the distal bipole of the ablation catheter at 5 mA and 2.0-ms pulse width duration, with ICE guidance, throughout the RV ventricular endocardium, along the papillary muscles, and along the MB. A pace-map match as high as 98% on both the Bard system and the CARTO-Paso system (Figures 3A–3C) was observed at the inferior-lateral free wall RV insertion site of the MB (Figure 3C), which was presumed to be the exit site of ventricular arrhythmia. Radiofrequency catheter ablation was performed at the best pace-match site, at the inferior-lateral free wall RV insertion site of the MB (Figures 3D and 3E and Online Supplemental Video 2). Radiofrequency energy up to 50 W was delivered at this site during sinus rhythm for up to 60–90 seconds during each ablation. To avoid complication while using 50 W power during ablation, we used a contact force sensing catheter to avoid any persistently high force application to avoid any potential risk of perforation. The impedance graph was also monitored throughout each ablation to evaluate for any rise that would be a precursor to a steam pop, and ablation was terminated if any sudden impedance rise was seen. In addition, the ablation catheter at the MB insertion site was constantly visualized using ICE during radiofrequency ablation. During initial ablation, salvos of PVCs and VT were observed that matched the morphology (>95% match) of the preablation PVC template. These salvos of PVCs resolved after further ablation of the RV MB lateral free wall insertion site. A total of 27 ablations were performed in the area surrounding the best pace-map match site near the RV MB insertion site. Postablation, reinduction of PVCs and VT was attempted with RV burst pacing and programmed electrical stimulation, both off and on isoproterenol infusion up to 5 µg/min. No ventricular ectopy was seen. We then reversed sedation completely and placed the patient in the left lateral decubitus position with all sheaths still in place, because of concern that the only preprocedural evidence of ventricular arrhythmia was in this position. In this left lateral decubitus position, no ventricular ectopy was seen. After a 30-minute waiting period without PVCs or VT, the procedure was terminated. Postablation, antiarrhythmic drugs were discontinued. Telemetry overnight did not show any clinical PVCs or NSVT, despite the patient lying in the left lateral decubitus position. Aspirin, atorvastatin, and an angiotensin-converting enzyme inhibitor were initiated and continued as an outpatient treatment because of newly diagnosed CAD. The patient was discharged home the next day in stable condition.

**Follow-up**

The patient resumed normal activity the day after the procedure and resumed work including performing surgery within 5 days after being discharged. He was evaluated in the outpatient arrhythmia clinic 1 month postablation and was noted to have no

Download English Version:

<https://daneshyari.com/en/article/8660682>

Download Persian Version:

<https://daneshyari.com/article/8660682>

[Daneshyari.com](https://daneshyari.com)