

Author's Accepted Manuscript

A simple method to ablate left-sided accessory pathways in a patient with coronary sinus ostial atresia and persistent left superior vena cava: A case report

Shohei Kataoka, Kenji Enta, Kyoichiro Yazaki, Mitsuru Kahata, Yasuhiro Ishii



PII: S2214-0271(16)30119-1
DOI: <http://dx.doi.org/10.1016/j.hrcr.2016.09.014>
Reference: HRCR304

To appear in: *HeartRhythm Case Reports*

Received date: 20 July 2016
Revised date: 17 September 2016
Accepted date: 19 September 2016

Cite this article as: Shohei Kataoka, Kenji Enta, Kyoichiro Yazaki, Mitsuru Kahata and Yasuhiro Ishii, A simple method to ablate left-sided accessory pathways in a patient with coronary sinus ostial atresia and persistent left superior vena cava: A case report, *HeartRhythm Case Reports*, <http://dx.doi.org/10.1016/j.hrcr.2016.09.014>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A simple method to ablate left-sided accessory pathways in a patient with coronary sinus ostial atresia and persistent left superior vena cava: A case report

Short title: Left-sided accessory pathways accompanied by coronary sinus ostial atresia and persistent left superior vena cava

Shohei Kataoka, MD¹, Kenji Enta, MD, PhD¹, Kyoichiro Yazaki, MD¹, Mitsuru Kahata, MD¹, Yasuhiro Ishii, MD, PhD¹

¹Department of Cardiology, Cardiovascular Center, Ogikubo Hospital

Corresponding author: Shohei Kataoka

Department of Cardiology, Cardiovascular Center, Ogikubo Hospital, 3-1-24 Imagawa, Suginami-ku, Tokyo 167-0035, Japan

Tel: +81-3-3399-1101, Fax: +81-3-3399-1107

E-mail: shoheikataoka0818@gmail.com

Key words: Wolff-Parkinson-White syndrome; Coronary sinus ostial atresia; Persistent left superior vena cava; Catheter ablation; Atrioventricular reentrant tachycardia

Conflict of interest: None

Introduction

Wolff-Parkinson-White (WPW) syndrome is characterized by a short PR interval, wide QRS complex, and paroxysmal supraventricular tachycardia.¹ An accessory atrioventricular connection is known to be the underlying mechanism of this syndrome. Accessory pathways

Download English Version:

<https://daneshyari.com/en/article/8660750>

Download Persian Version:

<https://daneshyari.com/article/8660750>

[Daneshyari.com](https://daneshyari.com)