


Accepted Manuscript

Prognostic impact of reducing myocardial ischemia identified using ECG-gated myocardial perfusion SPECT in Japanese patients with coronary artery disease: J-ACCESS 4 study

Mamoru Nanasato, Naoya Matsumoto, Kenichi Nakajima, Taishiro Chikamori, Masao Moroi, Kazuya Takehana, Mitsuru Momose, Hidetaka Nishina, Tokuo Kasai, Shunichi Yoda, Keisuke Kiso, Hiroyuki Yamamoto, Shigeyuki Nishimura, Akira Yamashina, Hideo Kusuoka, Atsushi Hirayama, Tsunehiko Nishimura



PII: S0167-5273(18)30218-3
DOI: doi:[10.1016/j.ijcard.2018.05.090](https://doi.org/10.1016/j.ijcard.2018.05.090)
Reference: IJCA 26503

To appear in:

Received date: 4 February 2018
Revised date: 12 May 2018
Accepted date: 23 May 2018

Please cite this article as: Mamoru Nanasato, Naoya Matsumoto, Kenichi Nakajima, Taishiro Chikamori, Masao Moroi, Kazuya Takehana, Mitsuru Momose, Hidetaka Nishina, Tokuo Kasai, Shunichi Yoda, Keisuke Kiso, Hiroyuki Yamamoto, Shigeyuki Nishimura, Akira Yamashina, Hideo Kusuoka, Atsushi Hirayama, Tsunehiko Nishimura, Prognostic impact of reducing myocardial ischemia identified using ECG-gated myocardial perfusion SPECT in Japanese patients with coronary artery disease: J-ACCESS 4 study. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Ijca(2017), doi:[10.1016/j.ijcard.2018.05.090](https://doi.org/10.1016/j.ijcard.2018.05.090)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Prognostic impact of reducing myocardial ischemia identified using ECG-gated
myocardial perfusion SPECT in Japanese patients with coronary artery disease:

J-ACCESS 4 study

Mamoru Nanasato^{a*}, Naoya Matsumoto^b, Kenichi Nakajima^c, Taishiro Chikamori^d,
Masao Moroi^e, Kazuya Takehana^f, Mitsuru Momose^g, Hidetaka Nishina^h, Tokuo Kasaiⁱ,
Shunichi Yoda^b, Keisuke Kiso^j, Hiroyuki Yamamoto^k, Shigeyuki Nishimura^l, Akira
Yamashina^d, Hideo Kusuoka^m, Atsushi Hirayama^b, Tsunehiko Nishimura^{n†}

^aCardiovascular Center, Nagoya Daini Red Cross Hospital, Nagoya, Japan

^bDivision of Cardiology, Department of Medicine, Nihon University School of
Medicine, Tokyo, Japan

^cDepartment of Nuclear Medicine, Kanazawa University Hospital, Kanazawa, Japan

^dDepartment of Cardiology, Tokyo Medical University, Tokyo, Japan

^eDivision of Cardiovascular Medicine, Toho University Ohashi Medical Center, Tokyo,
Japan

^fDepartment of Cardiology, Kansai Medical University, Hirakata, Japan

^gDepartment of Diagnostic Imaging and Nuclear Medicine, Tokyo Women's Medical
University, Tokyo, Japan

^hDepartment of Cardiology, Tsukuba Medical Center Hospital, Tsukuba, Japan

ⁱDepartment of Cardiology, Tokyo Medical University Hachioji Medical Center,
Hachioji, Japan.

^jDepartment of Radiology, National Cerebral and Cardiovascular Center, Suita, Japan

^kDepartment of Cardiology, Okayama Saiseikai General Hospital, Okayama, Japan

^lSaitama Medical University International Medical Center, Hidaka, Japan

^mNational Hospital Organization, Tokyo, Japan

ⁿGraduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto,
Japan

Running title: Ischemia reduction identified using MPI

*Correspondence during submission:

Download English Version:

<https://daneshyari.com/en/article/8661794>

Download Persian Version:

<https://daneshyari.com/article/8661794>

[Daneshyari.com](https://daneshyari.com)