


Accepted Manuscript

Coronary artery ectasia, an independent predictor of no-reflow after primary PCI for ST-elevation myocardial infarction

H.C.F. Schram, V.V. Hemradj, R.S. Hermanides, E. Kedhi, J.P. Ottervanger, the Zwolle Myocardial Infarction Study Group



PII: S0167-5273(18)31048-9
DOI: doi:[10.1016/j.ijcard.2018.04.120](https://doi.org/10.1016/j.ijcard.2018.04.120)
Reference: IJCA 26393

To appear in:

Received date: 13 February 2018
Revised date: 20 April 2018
Accepted date: 24 April 2018

Please cite this article as: H.C.F. Schram, V.V. Hemradj, R.S. Hermanides, E. Kedhi, J.P. Ottervanger, the Zwolle Myocardial Infarction Study Group, Coronary artery ectasia, an independent predictor of no-reflow after primary PCI for ST-elevation myocardial infarction. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Ijca*(2017), doi:[10.1016/j.ijcard.2018.04.120](https://doi.org/10.1016/j.ijcard.2018.04.120)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Coronary artery ectasia, an independent predictor of No-Reflow after Primary PCI for ST-Elevation Myocardial Infarction

H.C.F. Schram¹, MD, V.V. Hemradj², MD, R.S. Hermanides³, MD, PhD, E. Kedhi⁴, MD, PhD, J.P. Ottervanger⁵, MD, PhD, FESC, for the Zwolle Myocardial Infarction Study Group.

¹Isala, Zwolle, The Netherlands. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

²Isala, Zwolle, The Netherlands. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

³Isala, Zwolle, The Netherlands. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

⁴Isala, Zwolle, The Netherlands. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

⁵Isala, Zwolle, The Netherlands. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

Corresponding author:

Jan Paul Ottervanger, MD, PhD, FESC

Isala hospital, Department of Cardiology

Dr van Heesweg 2

8025 AB Zwolle, The Netherlands

Tel 31 38 4242374

E mail h.c.f.schram@isala.nl

No grant support was used for this study

Keywords: no-reflow, CAE, thrombus burden, primary percutaneous coronary intervention, ST-segment elevation myocardial infarction

Download English Version:

<https://daneshyari.com/en/article/8661871>

Download Persian Version:

<https://daneshyari.com/article/8661871>

[Daneshyari.com](https://daneshyari.com)