


Accepted Manuscript

Atrial fibrillation and ischemic events with rivaroxaban in patients with stable coronary artery disease (AFIRE): Protocol for a multicenter, prospective, randomized, open-label, parallel group study

Satoshi Yasuda, Koichi Kaikita, Hisao Ogawa, Masaharu Akao, Junya Ako, Tetsuya Matoba, Masato Nakamura, Katsumi Miyauchi, Nobuhisa Hagiwara, Kazuo Kimura, Atsushi Hirayama, Kunihiko Matsui



PII: S0167-5273(17)36755-4
DOI: doi:[10.1016/j.ijcard.2018.04.131](https://doi.org/10.1016/j.ijcard.2018.04.131)
Reference: IJCA 26404

To appear in:

Received date: 31 October 2017
Revised date: 12 April 2018
Accepted date: 30 April 2018

Please cite this article as: Satoshi Yasuda, Koichi Kaikita, Hisao Ogawa, Masaharu Akao, Junya Ako, Tetsuya Matoba, Masato Nakamura, Katsumi Miyauchi, Nobuhisa Hagiwara, Kazuo Kimura, Atsushi Hirayama, Kunihiko Matsui, Atrial fibrillation and ischemic events with rivaroxaban in patients with stable coronary artery disease (AFIRE): Protocol for a multicenter, prospective, randomized, open-label, parallel group study. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Ijca*(2017), doi:[10.1016/j.ijcard.2018.04.131](https://doi.org/10.1016/j.ijcard.2018.04.131)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Atrial Fibrillation and Ischemic events with Rivaroxaban in patiEnts with stable coronary artery disease (AFIRE): protocol for a multicenter, prospective, randomized, open-label, parallel group study

Satoshi Yasuda, MD, PhD¹, Koichi Kaikita, MD, PhD², Hisao Ogawa, MD, PhD³, Masaharu Akao, MD, PhD⁴, Junya Ako, MD, PhD⁵, Tetsuya Matoba, MD, PhD⁶, Masato Nakamura, MD, PhD⁷, Katsumi Miyauchi, MD, PhD⁸, Nobuhisa Hagiwara, MD, PhD⁹, Kazuo Kimura, MD, PhD¹⁰, Atsushi Hirayama, MD, PhD¹¹, Kunihiko Matsui, MD, PhD¹²

Affiliations

1. Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center Hospital, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
2. Department of Cardiovascular Medicine, Graduate School of Medical Sciences, Kumamoto University, 1-1-1 Honjo, Chuo-ku, Kumamoto 860-8556, Japan. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
3. National Cerebral and Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan. This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
4. Department of Cardiology, National Hospital Organization Kyoto Medical

Download English Version:

<https://daneshyari.com/en/article/8661892>

Download Persian Version:

<https://daneshyari.com/article/8661892>

[Daneshyari.com](https://daneshyari.com)