

## Accepted Manuscript

Morbidity and mortality after fragility hip fracture surgery in patients receiving vitamin K antagonists and direct oral anticoagulants

Tal Frenkel Rutenberg, Steven Velkes, Maria Vitenberg, Avi Leader, Yael Halavy, Pia Raanani, Mustafa Yassin, Galia Spectre



PII: S0049-3848(18)30327-X  
DOI: doi:[10.1016/j.thromres.2018.04.022](https://doi.org/10.1016/j.thromres.2018.04.022)  
Reference: TR 7016  
To appear in: *Thrombosis Research*  
Received date: 1 March 2018  
Revised date: 17 April 2018  
Accepted date: 20 April 2018

Please cite this article as: Tal Frenkel Rutenberg, Steven Velkes, Maria Vitenberg, Avi Leader, Yael Halavy, Pia Raanani, Mustafa Yassin, Galia Spectre , Morbidity and mortality after fragility hip fracture surgery in patients receiving vitamin K antagonists and direct oral anticoagulants. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Tr(2017), doi:[10.1016/j.thromres.2018.04.022](https://doi.org/10.1016/j.thromres.2018.04.022)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Morbidity and mortality after fragility hip fracture surgery in patients receiving vitamin K antagonists and direct oral anticoagulants

Tal Frenkel Rutenberg MD\*<sup>1</sup>, Steven Velkes MD\*<sup>1,4</sup>, Maria Vitenberg<sup>1</sup>, Avi Leader MD<sup>2,4</sup>, Yael Halavy<sup>2</sup>, Pia Raanani MD<sup>2,4</sup>, Mustafa Yassin<sup>3,4</sup>, Galia Spectre MD, PhD<sup>2,4</sup>

<sup>1</sup> Department of Orthopedics, Rabin Medical Center, Beilinson Hospital, Petah-Tikva, Israel.

<sup>2</sup> Institution of Hematology, Coagulation Unit, Rabin Medical Center, Petah-Tikva, Israel

<sup>3</sup> Department of Orthopedics, Hasharon Hospital, Rabin Medical Center, Petah-Tikva, Israel

<sup>4</sup> Sackler Faculty of Medicine, Tel-Aviv University, Tel Aviv, Israel.

\*These authors contributed equally to the study.

Corresponding author: Dr. Frenkel Rutenberg Tal, Orthopedic Department, Rabin Medical Center, Beilinson Hospital, 39 Jabotinsky St., Petah Tikva 49100.

Telephone: +972-54-4390190; e-mail: tal\_frenkel@hotmail.com

Funding: none

All authors declare no conflict of interests.

Word count: 4725

## ABSTRACT

**Introduction:** Early surgical treatment is recommended to reduce morbidity and mortality in patients with fragility hip fractures. Anticoagulation treatment poses a surgical challenge. While the action of vitamin K antagonists (VKAs) can be reversed, for direct oral anticoagulants (DOACs) antidote is only available for dabigatran. We aimed to assess the

Download English Version:

<https://daneshyari.com/en/article/8679381>

Download Persian Version:

<https://daneshyari.com/article/8679381>

[Daneshyari.com](https://daneshyari.com)