

Diagnostic Assessment & Prognosis

Phenoconversion from probable rapid eye movement sleep behavior disorder to mild cognitive impairment to dementia in a population-based sample

Youngsin Jung^a, Brendon P. Boot^b, Michelle M. Mielke^c, Tanis J. Ferman^d, Yonas E. Geda^{e,f}, Eric McDade^g, Teresa J. H. Christianson^c, David S. Knopman^a, Erik K. St Louis^{a,h}, Michael H. Silber^{a,h}, Ronald C. Petersen^a, Bradley F. Boeve^{a,h,*}

^aDepartment of Neurology, Mayo Clinic, Rochester, MN, USA

^bDepartment of Neurology, Brigham and Women's Hospital, Boston, MA, USA

^cDepartment of Health Sciences Research, Mayo Clinic, Rochester, MN, USA

^dDepartment of Psychiatry and Psychology, Mayo Clinic, Jacksonville, FL, USA

^eDepartment of Psychiatry and Psychology, Mayo Clinic, Scottsdale, AZ, USA

^fDepartment of Neurology, Mayo Clinic, Scottsdale, AZ, USA

^gDepartment of Neurology, Washington University School of Medicine in St. Louis, St. Louis, MO, USA

^hCenter for Sleep Medicine, Mayo Clinic, Rochester, MN, USA

Abstract

Introduction: Rapid eye movement sleep behavior disorder (RBD) is strongly associated with synucleinopathies. In 2012, we reported an increased risk of mild cognitive impairment (MCI) and Parkinson disease (PD) in cognitively normal Olmsted County, Minnesota, residents, aged 70 to 89 years with probable RBD. Here, we examine their progression to dementia and other neurodegenerative phenotypes.

Methods: Fifteen participants with RBD who were diagnosed with either MCI or PD were longitudinally followed, and their subsequent clinical courses were reviewed.

Results: Over 6.4 ± 2.9 years, six of the 14 participants with MCI developed additional neurodegenerative signs, five of whom had Lewy body disease features. Four of them progressed to dementia at a mean age 84.8 ± 4.9 years, three of whom met the criteria for probable dementia with Lewy bodies. One subject with PD developed MCI, but not dementia.

Discussion: Our findings from the population-based sample of Olmsted County, Minnesota, residents suggest that a substantial number of RBD patients tend to develop overt synucleinopathy features over time, and RBD patients who develop MCI and subsequent dementia have clinical features most consistent with dementia with Lewy bodies.

© 2017 The Authors. Published by Elsevier Inc. on behalf of the Alzheimer's Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords:

Rapid eye movement sleep behavior disorder; Mild cognitive impairment; Dementia with Lewy bodies; Parkinson disease; Synucleinopathy

1. Introduction

Rapid eye movement sleep behavior disorder (RBD) is a parasomnia characterized by complex motor behaviors (e.g., shouting, screaming, kicking, and punching) suggestive of

dream enactment [1]. The prevalence of RBD increases in older adults, especially in those with neurodegenerative conditions [2,3]. In particular, RBD is strongly associated with synucleinopathies, including Parkinson disease (PD), dementia with Lewy bodies (DLB), and multiple system atrophy [4,5]. Longitudinal cohort studies of patients with idiopathic RBD suggest that most of these patients ultimately develop an overt neurodegenerative phenotype

*Corresponding author. Tel.: +1 507-538-1038; Fax: +1 507-538-6012.

E-mail address: bboeve@mayo.edu

with the rate of phenoconversion as high as 91% at 14 years [6–8]. However, these studies are based on sleep center cohorts in which referral bias toward more severe disease is likely to be present, and therefore may not reflect findings in the general population.

To assess population-based risk of neurodegenerative disease in RBD, we previously studied cognitively normal individuals aged 70 to 89 years with probable RBD (pRBD) in Olmsted County, Minnesota, and reported a 2.2-fold-increased risk of developing mild cognitive impairment (MCI) or PD over 4 years [3]. Since the initial report, we have longitudinally followed the participants with pRBD who developed MCI or PD as part of the Mayo Clinic Study of Aging (MCSA), a longitudinal population-based study of aging [9]. Here, we examine their progression to dementia and other neurodegenerative phenotypes.

2. Methods

The details of subject recruitment and diagnostic evaluation are described in our previous report [3]. Briefly, the original cohort included the MCSA participants aged 70 to 89 years who were randomly selected residents of Olmsted County, Minnesota, using the medical records linkage system of the Rochester Epidemiology Project. All individuals in the original cohort were cognitively normal without PD at the time of their initial enrollment in the MCSA. These participants were longitudinally followed based on a standardized protocol [9]. The diagnosis of pRBD was made using the Mayo Sleep Questionnaire [10]. A consensus diagnosis of MCI, dementia, or PD was made according to our previously published criteria [9].

A total of 15 participants with pRBD who were diagnosed with either MCI or PD at the time of our original analysis

were included in the present study. Follow-up and autopsy data were obtained from the MCSA and/or review of their medical records.

Informed consent was obtained from the participants and their family members or significant others. All study protocols were approved by the Mayo Clinic and Olmsted Medical Center institutional review boards.

3. Results

The demographic and clinical characteristics of the participants are summarized in the Table 1. The participants were followed for an average 6.4 ± 2.9 years after their MCI or PD diagnosis. Of the 14 participants with MCI, four with amnesic MCI developed dementia at a mean age 84.8 ± 4.9 years: three met the criteria for probable DLB [11] and one for Alzheimer disease dementia [12]. The estimated onset of dementia began 13.8 ± 6.2 years after pRBD diagnosis and 4.3 ± 2.8 years after MCI diagnosis. All participants with probable DLB had well-formed visual hallucinations, and one of them also had parkinsonism. One participant with nonamnesic MCI developed visual hallucinations 2 years after the MCI diagnosis and another developed parkinsonism 4 years after the MCI diagnosis, but neither met criteria for dementia. One participant with PD subsequently developed nonamnesic multi-domain MCI 10 years after the PD diagnosis. The remainder of the participants continued to have MCI until their death or last follow-up (5.9 ± 2.9 years), except for one who was reclassified as having normal cognition 7 years before her death and remained cognitively normal until the death. Her prior cognitive insufficiency was presumed related to suboptimally controlled depression and/or polypharmacy. This individual underwent autopsy and

Table 1
Demographic and clinical characteristics of the updated probable RBD cohort

Subject	Sex	Age at the onset of RBD (year)	Age at the onset of MCI (year)	MCI type	Age at the onset of dementia (year)	Dementia type	Age at death (year)	Age at last follow-up (year)	Other features
1	Male	71	77	A	84	Probable DLB	85	n/a	VH, parkinsonism
2	Male	63	84	A	85	Probable DLB	89	n/a	VH
3	Female	78	85	A	91	Probable DLB	n/a	92	VH
4	Male	72	76	A	79	Probable AD	81	n/a	
5	Female	32	82*	n/a	n/a	n/a	91	n/a	
6	Male	66	87	A	n/a	n/a	88	n/a	
7	Male	23	85	A	n/a	n/a	n/a	93	
8	Male	82	84	A	n/a	n/a	n/a	93	
9	Male	71	81	A, M	n/a	n/a	85	n/a	
10	Male	69	74	A, M	n/a	n/a	n/a	82	
11	Female	84	86	NA	n/a	n/a	89	n/a	VH
12	Male	27	78	NA	n/a	n/a	84	n/a	Parkinsonism
13	Male	30	78	NA	n/a	n/a	n/a	86	
14	Male	47	76	NA, M	n/a	n/a	78	n/a	PD
15	Male	34	84	U	n/a	n/a	87	n/a	

Abbreviations: A, amnesic; AD, Alzheimer disease; DLB, dementia with Lewy bodies; M, multidomain; MCI, mild cognitive impairment; n/a, not applicable; NA, nonamnesic; PD, Parkinson disease; RBD, rapid eye movement sleep behavior disorder; U, unknown type; VH, visual hallucinations.

*Classification changed from amnesic MCI to normal 7 years before her death.

Download English Version:

<https://daneshyari.com/en/article/8680331>

Download Persian Version:

<https://daneshyari.com/article/8680331>

[Daneshyari.com](https://daneshyari.com)