Accepted Manuscript



Non-motor outcomes of subthalamic stimulation in Parkinson's disease depend on location of active contacts

Haidar Salimi Dafsari, Jan Niklas Petry-Schmelzer, K. Ray-Chaudhuri, Keyoumars Ashkan, Luca Weis, Till A. Dembek, Michael Samuel, Alexandra Rizos, Monty Silverdale, Michael T. Barbe, Gereon R. Fink, Julian Evans, Pablo Martinez-Martin, Angelo Antonini, Veerle Visser-Vandewalle, Lars Timmermann

PII: S1935-861X(18)30092-5

DOI: 10.1016/j.brs.2018.03.009

Reference: BRS 1217

To appear in: *Brain Stimulation*

Received Date: 24 July 2017

Revised Date: 6 March 2018

Accepted Date: 12 March 2018

Please cite this article as: Dafsari HS, Petry-Schmelzer JN, Ray-Chaudhuri K, Ashkan K, Weis L, Dembek TA, Samuel M, Rizos A, Silverdale M, Barbe MT, Fink GR, Evans J, Martinez-Martin P, Antonini A, Visser-Vandewalle V, Timmermann L, On behalf of EUROPAR, On behalf of the IPMDS Non Motor PD Study Group, Non-motor outcomes of subthalamic stimulation in Parkinson's disease depend on location of active contacts, *Brain Stimulation* (2018), doi: 10.1016/j.brs.2018.03.009.

Brain Stimulation (Original Article)

Non-motor outcomes of subthalamic stimulation in Parkinson's disease depend on location of active contacts

Haidar Salimi Dafsari[1, 2]*, MD, Jan Niklas Petry-Schmelzer[1]*, MD, K. Ray-Chaudhuri[2, 3], Prof., Keyoumars Ashkan[2], Prof., Luca Weis[4], MD, Till A. Dembek[1], MD, Michael Samuel[2], MD/PhD, Alexandra Rizos[2], MSc, Monty Silverdale[5], MD/PhD, Michael T. Barbe[1], MD, Gereon R. Fink[1, 6], Prof., Julian Evans[4], Prof., Pablo Martinez-Martin[7], Prof., Angelo Antonini[4], Prof., Veerle Visser-Vandewalle[8], Prof., Lars Timmermann[1, 9], Prof.,
On behalf of EUROPAR and the IPMDS Non Motor PD Study Group

[1] Department of Neurology, University Hospital Cologne, Cologne, Germany.
[2] National Parkinson Foundation International Centre of Excellence, King's College Hospital, London, United Kingdom.
[3] The Maurice Wohl Clinical Neuroscience Institute, King's College London, London, United Kingdom.
[4] Department of Neurology, IRCCS, San Camillo, Venice, Italy.
[5] Department of Neurology and Neurosurgery, Salford Royal Foundation Trust, Manchester Academic Health Science Centre, University of Manchester, Greater Manchester, United Kingdom.
[6] Cognitive Neuroscience, Institute of Neuroscience and Medicine (INM-3), Research Center Jülich, Jülich, Germany.
[7] National Center of Epidemiology and CIBERNED, Carlos III Institute of Health, Madrid, Spain.
[8] Department of Stereotaxy and Functional Neurosurgery, University Hospital Cologne, Cologne, Germany.
[9] Department of Neurology, University Hospital Giessen and Marburg, Campus Marburg, Germany.

Character count title: 106
Number of references: 41
Number of tables: 3
Number of figures: 2
Word count abstract: 250/250
Word count text: 3169/4000
Supplemental Data: 1 (Supplemental table e-1)

Corresponding Author:
Dr. Haidar S. Dafsari
Department of Neurology
University Hospital Cologne
Kerpenerstr. 62
50924 Cologne, Germany
Tel.: 0221-478-4014
E-mail: haidar.dafsari@uk-koeln.de

Statistical analysis conducted by Dr. Haidar S. Dafsari, MD, University Hospital Cologne, Germany, and National Parkinson Foundation International Centre of Excellence, King's College Hospital, London, United Kingdom.

Search terms: Deep brain stimulation; subthalamic nucleus; non-motor symptoms; non motor symptoms; quality of life

* These authors contributed equally to the manuscript.