


Accepted Manuscript

Title: Postural control before and after cerebrospinal fluid shunt surgery in idiopathic normal pressure hydrocephalus

Authors: Yasutaka Nikaido, Toshihiro Akisue, Hideyuki Urakami, Yoshinaga Kajimoto, Kenji Kuroda, Yuki Kawami, Hisatomo Sato, Yoshiyuki Ohta, Tetsuya Hinoshita, Yuka Iwai, Tadayuki Nishiguchi, Hiroshi Ohno, Ryuichi Saura



PII: S0303-8467(18)30251-8
DOI: <https://doi.org/10.1016/j.clineuro.2018.06.032>
Reference: CLINEU 5075

To appear in: *Clinical Neurology and Neurosurgery*

Received date: 25-4-2018
Revised date: 17-6-2018
Accepted date: 29-6-2018

Please cite this article as: Nikaido Y, Akisue T, Urakami H, Kajimoto Y, Kuroda K, Kawami Y, Sato H, Ohta Y, Hinoshita T, Iwai Y, Nishiguchi T, Ohno H, Saura R, Postural control before and after cerebrospinal fluid shunt surgery in idiopathic normal pressure hydrocephalus, *Clinical Neurology and Neurosurgery* (2018), <https://doi.org/10.1016/j.clineuro.2018.06.032>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title:

Postural control before and after cerebrospinal fluid shunt surgery in idiopathic normal pressure hydrocephalus

Authors:

Yasutaka Nikaido^{a,b,*}, Toshihiro Akisue^b, Hideyuki Urakami^a, Yoshinaga Kajimoto^c, Kenji Kuroda^a, Yuki Kawami^{b,d}, Hisatomo Sato^a, Yoshiyuki Ohta^a, Tetsuya Hinoshita^a, Yuka Iwai^a, Tadayuki Nishiguchi^a, Hiroshi Ohno^a, Ryuichi Saura^e

Affiliations:

- a. Clinical department of Rehabilitation, Osaka Medical College Hospital, Osaka, Japan
- b. Department of Rehabilitation Science, Graduate School of Health Sciences, Kobe University, Kobe, Japan
- c. Department of Neurosurgery, Division of Surgery, Osaka Medical College, Osaka, Japan
- d. Department of Rehabilitation, Hyogo Rehabilitation Centre at Nishiharima, Hyogo, Japan
- e. Department of Physical and Rehabilitation Medicine, Division of Comprehensive Medicine, Osaka Medical College, Osaka, Japan

Corresponding Author:

Yasutaka Nikaido

Clinical department of Rehabilitation, Osaka Medical College Hospital, 2-7, Daigaku-machi, Takatsuki, Osaka 569-8686, Japan.

Tel.: +81-72-683-1221.

E-mail address: reh055@osaka-med.ac.jp

Highlights

- Postural sway was greater in iNPH patients before shunt surgery.
- Voluntary COP movements improved in iNPH patients at 1 week post-surgery.
- TUG value and voluntary COP movements were correlated in iNPH patients.
- No correlation observed between TUG value and quiescent standing.
- CSF shunting in iNPH patients may improve voluntary postural control.

Download English Version:

<https://daneshyari.com/en/article/8681650>

Download Persian Version:

<https://daneshyari.com/article/8681650>

[Daneshyari.com](https://daneshyari.com)