


Accepted Manuscript

Title: Surgical approach for suprasellar hemangioblastomas preserving the pituitary stalk: review of the literature and report of a further case

Authors: Nabeel Alshafai, Rodolfo Maduri, Mrigank Shail, Domenico Chirchiglia, David Meyronet, Francesco Signorelli



PII: S0303-8467(18)30113-6
DOI: <https://doi.org/10.1016/j.clineuro.2018.03.010>
Reference: CLINEU 4965

To appear in: *Clinical Neurology and Neurosurgery*

Received date: 23-10-2017
Revised date: 3-3-2018
Accepted date: 8-3-2018

Please cite this article as: Alshafai N, Maduri R, Shail M, Chirchiglia D, Meyronet D, Signorelli F, Surgical approach for suprasellar hemangioblastomas preserving the pituitary stalk: review of the literature and report of a further case, *Clinical Neurology and Neurosurgery* (2010), <https://doi.org/10.1016/j.clineuro.2018.03.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Surgical approach for suprasellar hemangioblastomas preserving the pituitary stalk: review of the literature and report of a further case

Nabeel Alshafai¹, Rodolfo Maduri², Mrigank Shail³, Domenico Chirchiglia⁴, David Meyronet⁵,
and Francesco Signorelli^{6,7}

¹Alshafai Neurosurgical Academy (A.N.A.), Toronto, Ontario, Canada

²Department of Clinical Neurosciences, Service of Neurosurgery, Centre Hospitalier
Universitaire Vaudois, Lausanne, Switzerland

³Xavier University School of Medicine, Aruba, The Netherlands

⁴Department of Medical Sciences, University « Magna Græcia » of Catanzaro, Italy;

⁵Centre de Pathologie et de Neuropathologie Est, Hospices Civils de Lyon, Lyon, France

⁶Department of Neurosurgery, Hôpital Pierre Wertheimer, Hospices Civils de Lyon, Lyon,
France

⁷Division of Neurosurgery, Department of Basic Medical Sciences, Neurosciences and Sense
Organs, University “Aldo Moro” of Bari, Italy.

Corresponding Author: Francesco Signorelli, MD, Msc, Division of Neurosurgery, Department
of Basic Medical Sciences, Neurosciences and Sense Organs, University “Aldo Moro” of Bari,
Piazza Giulio Cesare, 11, 70124 Bari, Italy.

Email: signorelli2007@gmail.com

Presented in part in abstract form and oral presentation form at the 63rd National Congress of the
Italian Society of Neurosurgery, September 25-27, 2014, Vicenza, Italy.

Highlights

- Pituitary stalk HBLs on MRI avidly enhance and are associated with flow voids
- Surgery is indicated for symptomatic lesions
- Radical removal with stalk sparing is the treatment of choice
- Surgery is challenging due to the location and hypervascular nature of the HBL
- OZ approach with EAC improves outcome compared to subfrontal and pterional approach

Download English Version:

<https://daneshyari.com/en/article/8681827>

Download Persian Version:

<https://daneshyari.com/article/8681827>

[Daneshyari.com](https://daneshyari.com)