Accepted Manuscript

Title: Epidemiology and Short-term Surgical Outcomes of Children Presenting with Cerebellar Tumors

Authors: Balagangadhar R. Totapally, Ashish H. Shah, Toba

Niazi

PII: S0303-8467(18)30089-1

DOI: https://doi.org/10.1016/j.clineuro.2018.02.038


Reference: CLINEU 4949

To appear in: Clinical Neurology and Neurosurgery

Received date: 19-12-2017 Revised date: 14-2-2018 Accepted date: 25-2-2018

Please cite this article as: Totapally BR, Shah AH, Niazi T, Epidemiology and Short-term Surgical Outcomes of Children Presenting with Cerebellar Tumors, *Clinical Neurology and Neurosurgery* (2010), https://doi.org/10.1016/j.clineuro.2018.02.038

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Totapally et al. 1

Epidemiology and Short-term Surgical Outcomes of Children Presenting with Cerebellar

Tumors

Short Title: Surgical outcomes of cerebellar tumors

Balagangadhar R. Totapally, MD, 1,2 Ashish H. Shah, MD, 3,4 Toba Niazi, MD, 2,3,4

Affiliations: ¹Division of Critical Care Medicine, Nicklaus Children's Hospital, Miami, FL 33155;

²Herbert Wertheim College of Medicine, Florida International University, Miami, FL 33199;

³Division of Neurosurgery, Nicklaus Children's Hospital, Miami, FL 33155, ⁴ Department of

Neurosurgery, University of Miami Miller School of Medicine, Miami, FL 33133

Address for Correspondence: Balagangadhar R. Totapally, MD, Division of Critical Care

Medicine, Nicklaus Children's Hospital, 3100 SW 62nd Avenue, Miami, FL 33165.

E-mail: bala.totapally@mch.com; Tel: 305 662 2639; Fax: 305 663 0530

Highlights:

The prevalence of children with cerebellar tumors is 1.7 per 10,000 discharges.

Many children (37%) presenting with cerebellar tumors required EVD and/or VP shunt.

The hospital mortality is low (<1%) in children presenting with cerebellar tumors.

12% of these children are discharged to skilled nursing facilities.

Abstract

Objective: Posterior fossa tumor surgery in children poses a significant morbidity and mortality.

Large multi-institutional datasets characterizing the epidemiology and morbidity of children

1

Download English Version:

https://daneshyari.com/en/article/8681848

Download Persian Version:

https://daneshyari.com/article/8681848

<u>Daneshyari.com</u>