Accepted Manuscript

Title: Spontaneous speech in patients with gliomas in eloquent

areas: Evaluation until 1 year after surgery

Authors: Djaina Satoer, Arnaud Vincent, Leonie Ruhaak, Marion Smits, Clemens Dirven, Evy Visch-Brink

PII: \$0303-8467(18)30063-5

DOI: https://doi.org/10.1016/j.clineuro.2018.02.018


Reference: CLINEU 4929

To appear in: Clinical Neurology and Neurosurgery

Received date: 11-1-2018 Revised date: 8-2-2018 Accepted date: 11-2-2018

Please cite this article as: Satoer D, Vincent A, Ruhaak L, Smits M, Dirven C, Visch-Brink E, Spontaneous speech in patients with gliomas in eloquent areas: Evaluation until 1 year after surgery, *Clinical Neurology and Neurosurgery* (2010), https://doi.org/10.1016/j.clineuro.2018.02.018

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

TITLE PAGE

Title of the paper: Spontaneous speech in patients with gliomas in eloquent areas: evaluation until 1 year after surgery.

Names of the authors: Djaina Satoer^a PhD, Arnaud Vincent^a MD PhD, Leonie Ruhaak^a, MSc, Marion Smits^b MD PhD, Clemens Dirven^a MD PhD, Evy Visch-Brink^a PhD

Affiliations: Departments of ^aNeurosurgery and ^bRadiology and Nuclear Medicine Erasmus MC - University Medical Center Rotterdam, Rotterdam, The Netherlands.

Corresponding author: Djaina Satoer, e-mail: d.satoer@erasmusmc.nl.

Present address: Department of Neurosurgery, Erasmus MC - University Medical Center, room BA 479, P.O. Box 2040, 3000 CA Rotterdam, telephone +31 010 7034077.

Declaration of interest: none

HIGHLIGHTS

- Glioma patients have impaired spontaneous speech with naming and fluency deficits.
- Spontaneous speech analysis is sensitive to detect long-term language decline.
- Most spontaneous speech variables did not correlate with standard language tests.
- Spontaneous speech has added value to standard tests for diagnosis of deficits.

ABSTRACT

Objective: Glioma patients often complain about problems in daily conversation with a negative impact on quality of life. Disorders in standardized language tests (e.g. naming and fluency), are frequently observed. Most studies claim recovery of language functions within 3 months. However, long-term effects of surgery on spontaneous speech remain unknown.

Patients and Methods: Eighteen glioma patients were compared to healthy controls in spontaneous speech variables: Type Token Ratio (TTR), Mean Length of Utterance words (MLUw), Incomplete Sentences, Self-corrections and Repetitions. Boston Naming Test (BNT) and Category Fluency (CF) were also assessed. We compared: pre- and 3 months postoperatively (T1-T2), 3 months and 1 year postoperatively (T2-T3), pre- and 1 year postoperatively (T1-T3). Correlations were computed between deviating variables and BNT/CF, tumor localization, and tumor grade.

Results: Patients had deficits in Incomplete sentences (T1, T2, T3), TTR (T2,T3), MLUw (T3) and Self-corrections (T2). Between T1-T2 no decline was present. Between T2-T3 and T1-T3, there was a decrease of MLUw, Self-corrections and Repetitions and an increase of

Download English Version:

https://daneshyari.com/en/article/8681899

Download Persian Version:

https://daneshyari.com/article/8681899

Daneshyari.com