

## Accepted Manuscript

Diagnostic accuracy of quantitative EEG to detect delayed cerebral ischemia after subarachnoid hemorrhage: A preliminary study

B. Balança, F. Dailler, S. Boulogne, T. Ritzenthaler, F. Gobert, S. Rheims, N. Andre-Obadia

PII: S1388-2457(18)31133-7

DOI: <https://doi.org/10.1016/j.clinph.2018.06.013>

Reference: CLINPH 2008578

To appear in: *Clinical Neurophysiology*


Received Date: 18 December 2017

Revised Date: 24 April 2018

Accepted Date: 15 June 2018

Please cite this article as: Balança, B., Dailler, F., Boulogne, S., Ritzenthaler, T., Gobert, F., Rheims, S., Andre-Obadia, N., Diagnostic accuracy of quantitative EEG to detect delayed cerebral ischemia after subarachnoid hemorrhage: A preliminary study, *Clinical Neurophysiology* (2018), doi: <https://doi.org/10.1016/j.clinph.2018.06.013>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Diagnostic accuracy of quantitative EEG to detect delayed cerebral ischemia after subarachnoid hemorrhage: a preliminary study<sup>†</sup>**

Balança B<sup>1,2,3\*</sup>, Dailier F<sup>1</sup>, Boulogne S<sup>2,4</sup>, Ritzenthaler T<sup>1,5</sup>, Gobert F<sup>1,6</sup>, Rheims S<sup>2,4</sup>, Andre-Obadia N<sup>4,7</sup>.

<sup>1</sup> Hospices Civils de Lyon, Hôpital Pierre Wertheimer, Department of anesthesiology and intensive care medicine, Lyon, France

<sup>2</sup> Inserm U1028, CNRS UMR 5292, Lyon Neuroscience Research Centre, Team TIGER, Lyon, France

<sup>3</sup> Université Claude Bernard Lyon 1, Centre Lyonnais d'Enseignement par la Simulation en Santé, SAMSEI, Lyon, France

<sup>4</sup> Hospices Civils de Lyon, Hôpital Pierre Wertheimer, Department of Functional Neurology and Epileptology, Lyon, France

<sup>5</sup> CREATIS, CNRS UMR 5220INSERM U1044, Université Lyon 1, Villeurbanne, France

<sup>6</sup> Inserm U1028, CNRS UMR 5292, Lyon Neuroscience Research Centre, Team IMAPCT, Lyon, France

<sup>7</sup> Inserm U1028, CNRS UMR 5292, Lyon Neuroscience Research Centre, Team NEUROPAIN, Lyon, France

\* **Corresponding author:** Dr Baptiste Balança, Department of neurologic anesthesiology and intensive care medicine, Hôpital Pierre Wertheimer, Hospices Civils de Lyon, Lyon, France.

Tel: +33 6 23 91 05 94

mail: [baptiste.balanca@gmail.com](mailto:baptiste.balanca@gmail.com)

<sup>†</sup> This study was performed in the Department of anesthesiology and intensive care medicine of the Pierre Wertheimer Hospital, Hospices Civils de Lyon, Lyon, France.

Download English Version:

<https://daneshyari.com/en/article/8682166>

Download Persian Version:

<https://daneshyari.com/article/8682166>

[Daneshyari.com](https://daneshyari.com)