

## Accepted Manuscript

Title: Clinical and Electrographic Features of Sunflower Syndrome

Authors: Fiona M. Baumer, Brenda E. Porter

PII: S0920-1211(17)30618-6

DOI: <https://doi.org/10.1016/j.eplepsyres.2018.03.002>

Reference: EPIRES 5915

To appear in: *Epilepsy Research*

Received date: 5-12-2017

Revised date: 26-2-2018

Accepted date: 1-3-2018



Please cite this article as: Baumer, Fiona M., Porter, Brenda E., Clinical and Electrographic Features of Sunflower Syndrome. *Epilepsy Research* <https://doi.org/10.1016/j.eplepsyres.2018.03.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Clinical and Electrographic Features of Sunflower Syndrome

Fiona M. Baumer<sup>a</sup> MD and Brenda E. Porter<sup>a</sup> MD PhD

<sup>a</sup>Stanford University School of Medicine, Department of Neurology, Division of Child Neurology

### Corresponding Author:

Fiona M. Baumer, MD  
Lucile Packard Children's Hospital  
Stanford University Medical School  
750 Welch, Suite 317  
Palo Alto, CA 94304  
T: (650) 736-0885  
F: (650) 723-7299  
[fbaumer@stanford.edu](mailto:fbaumer@stanford.edu)

### Other Authors:

Brenda Porter, MD, PhD  
Same postal address as corresponding author.  
[brenda2@stanford.edu](mailto:brenda2@stanford.edu)

Title:	58 characters (with spaces)
Abstract:	287 words
Body:	2696 words
Number of tables:	3
Number of figures:	0
Number of references:	21

**Financial Disclosures/Declarations of Interest:** None

### Highlights

- Self-induced seizures in photosensitive patients are difficult to treat.
- Typical patients are school-aged and often have cognitive/attention difficulties.
- Seizures consist of eyelid myoclonia +/- absence; 50% have spontaneous seizures.
- Consider valproate or polypharmacy early; avoid levetiracetam monotherapy.
- Counsel on trigger avoidance. Measure treatment response with EEGs done outside.

Download English Version:

<https://daneshyari.com/en/article/8684147>

Download Persian Version:

<https://daneshyari.com/article/8684147>

[Daneshyari.com](https://daneshyari.com)