

Regenerating islet-derived 1α (REG-1α) protein increases tau phosphorylation in cell and animal models of tauopathies

Mireille Moussaed, Sylvaine Huc-Brandt, Nicolas Cubedo, Michele Silhol, Samy Murat, Marie-Christine Lebart, Gabor Kovacs, Jean-Michel Verdier, Francoise Trousse, Mireille Rossel, Anne Marcilhac

PII: S0969-9961(18)30351-6
DOI: doi:10.1016/j.nbd.2018.07.029
Reference: YNBDI 4239

To appear in: *Neurobiology of Disease*

Received date: 21 February 2018
Revised date: 15 June 2018
Accepted date: 28 July 2018

Please cite this article as: Mireille Moussaed, Sylvaine Huc-Brandt, Nicolas Cubedo, Michele Silhol, Samy Murat, Marie-Christine Lebart, Gabor Kovacs, Jean-Michel Verdier, Francoise Trousse, Mireille Rossel, Anne Marcilhac , Regenerating islet-derived 1α (REG-1α) protein increases tau phosphorylation in cell and animal models of tauopathies. Ynbdi (2018), doi:10.1016/j.nbd.2018.07.029

Regenerating islet-derived 1α (REG-1α) protein increases tau phosphorylation in cell and animal models of tauopathies.

Mireille Moussaed.[1], Sylvaine Huc-Brandt [1], Nicolas Cubedo [1], Michele Silhol [1], Samy Murat[1], Marie-Christine Lebart [1], Gabor Kovacs [2], Jean-Michel Verdier [1], Francoise Trousse [1], Mireille Rossel [1a], Anne Marcilhac [1a]*

[1]MMDN, Univ. Montpellier, EPHE, INSERM, U1198, PSL University, Montpellier, F-34095

France.

[2]Institute of Neurology, Neurodegeneration Research Group, Medical University of Vienna, Vienna, Austria.

[a] Both authors contributed equally to this work.

*Corresponding author:

Dr Marcilhac Anne

INSERM U1198, Montpellier University, Place E. Bataillon, CC 105, 34095 Montpellier cedex

05, France.

anne.marcilhac@umontpellier.fr