

Accepted Manuscript

Can anomalous diffusion models in magnetic resonance imaging be used to characterise white matter tissue microstructure?

Qiang Yu, David Reutens, Viktor Vegh

PII: S1053-8119(18)30262-3

DOI: [10.1016/j.neuroimage.2018.03.052](https://doi.org/10.1016/j.neuroimage.2018.03.052)

Reference: YNIMG 14821

To appear in: *NeuroImage*


Received Date: 4 December 2017

Revised Date: 13 March 2018

Accepted Date: 22 March 2018

Please cite this article as: Yu, Q., Reutens, D., Vegh, V., Can anomalous diffusion models in magnetic resonance imaging be used to characterise white matter tissue microstructure?, *NeuroImage* (2018), doi: [10.1016/j.neuroimage.2018.03.052](https://doi.org/10.1016/j.neuroimage.2018.03.052).

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Can anomalous diffusion models in magnetic resonance imaging be used to characterise
white matter tissue microstructure?**

Qiang Yu, David Reutens and Viktor Vegh*

Centre for Advanced Imaging, the University of Queensland, Brisbane, Queensland, Australia

Short running title: Probing of tissue microstructure via anomalous diffusion models

*Corresponding author:

Viktor Vegh

Centre for Advanced Imaging

Building 57

University of Queensland

Brisbane, QLD, 4072

Australia

E-mail: v.vegh@uq.edu.au

Phone: +61 7 3365 4100

Fax: +61 7 3346 0330

Word count for body of text: 8335

Download English Version:

<https://daneshyari.com/en/article/8686869>

Download Persian Version:

<https://daneshyari.com/article/8686869>

[Daneshyari.com](https://daneshyari.com)