Aberrant hemodynamic responses in autism: Implications for resting state fMRI functional connectivity studies

Wenjing Yan, D. Rangaprakash, Gopikrishna Deshpande

PII: S2213-1582(18)30122-0
DOI: doi:10.1016/j.nicl.2018.04.013
Reference: YNICL 1374

To appear in: *NeuroImage: Clinical*

Received date: 10 October 2017
Revised date: 28 March 2018
Accepted date: 11 April 2018

Please cite this article as: Wenjing Yan, D. Rangaprakash, Gopikrishna Deshpande , Aberrant hemodynamic responses in autism: Implications for resting state fMRI functional connectivity studies. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Ynicl(2017), doi:10.1016/j.nicl.2018.04.013

Aberrant Hemodynamic Responses in Autism: Implications for Resting State fMRI Functional Connectivity Studies

Wenjing Yan[1], D Rangaprakash[1,2], Gopikrishna Deshpande[1,3,4,5*]

[1]AU MRI Research Center, Department of Electrical and Computer Engineering, Auburn University, Auburn, AL, USA

[2]Department of Psychiatry and Biobehavioral Sciences, University of California Los Angeles, Los Angeles, CA, USA

[3]Department of Psychology, Auburn University, Auburn, AL, USA

[4] Center for Health Ecology and Equity Research, Auburn University, Auburn, AL, USA

[5]Alabama Advanced Imaging Consortium, Auburn University and University of Alabama Birmingham, AL, USA

***Correspondence to:**

Gopikrishna Deshpande, Ph.D.

AU MRI research center, Dept. of Electrical & Computer Engineering,

560 Devall Dr, Suite 266D,

Auburn University, Auburn, AL 36849, USA.

Tel: +1-334-844-7653; Fax: +1-334-844-0214; Email: gopi@auburn.edu

Disclosures: The authors report no competing interests.

Keywords: Resting-state fMRI, Deconvolution, Autism, Hemodynamic Response Function (HRF), HRF variability, Seed-based Functional Connectivity