

Accepted Manuscript

Hierarchical disruption in the Bayesian brain: Focal epilepsy and brain networks

Amir Omidvarnia, Mangor Pedersen, Richard E. Rosch, Karl J. Friston, Graeme D. Jackson



PII: S2213-1582(17)30125-0
DOI: doi: [10.1016/j.nicl.2017.05.019](https://doi.org/10.1016/j.nicl.2017.05.019)
Reference: YNICL 1034

To appear in: *NeuroImage: Clinical*

Received date: 21 December 2016
Revised date: 22 March 2017
Accepted date: 25 May 2017

Please cite this article as: Amir Omidvarnia, Mangor Pedersen, Richard E. Rosch, Karl J. Friston, Graeme D. Jackson, Hierarchical disruption in the Bayesian brain: Focal epilepsy and brain networks, *NeuroImage: Clinical* (2017), doi: [10.1016/j.nicl.2017.05.019](https://doi.org/10.1016/j.nicl.2017.05.019)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Hierarchical disruption in the Bayesian brain: Focal epilepsy and brain networks

Amir Omidvarnia¹, Mangor Pedersen¹, Richard E. Rosch², Karl J. Friston², Graeme D. Jackson^{1,3}

1. The Florey Institute of Neuroscience and Mental Health and University of Melbourne, Austin Campus, Heidelberg, VIC, Australia.
2. The Wellcome Trust Centre for Neuroimaging, Institute of Neurology, University College London, London, UK
3. Department of Neurology, Austin Health, Heidelberg, VIC, Australia.

Running title: Focal epilepsy and false inference in brain networks

Abstract

In this opinion paper, we describe a combined view of functional and effective brain connectivity along with the free-energy principle for investigating persistent disruptions in brain networks of patients with focal epilepsy. These changes are likely reflected in effective connectivity along the cortical hierarchy and construct the basis of increased functional connectivity in focal epilepsy. We propose a testable framework based on dynamic causal modelling and functional connectivity analysis with the capacity of explaining commonly observed connectivity changes during inter-ictal periods. We then hypothesise their possible relation with disrupted free-energy minimisation in the Bayesian brain. This may offer a new approach for neuroimaging to specifically develop and address hypotheses regarding the network pathomechanisms underlying epileptic phenotypes.

Keywords: focal epilepsy, free-energy principle, functional connectivity, effective connectivity, Bayesian inference, predictive coding

1. Background

Epilepsy is a dynamic disorder of the brain that is characterised by both paroxysmal abnormal states (i.e. epileptic seizures), and more persistent abnormalities across the functional brain networks (Powell et al., 2007). Advances in understanding the relationship between these observable phenomena and underlying pathophysiology have directly informed current treatment approaches, such as epilepsy surgery, and can potentially improve patient outcomes (Goodfellow et al., 2016). However, understanding the neuronal

Download English Version:

<https://daneshyari.com/en/article/8688520>

Download Persian Version:

<https://daneshyari.com/article/8688520>

[Daneshyari.com](https://daneshyari.com)