

Parietal operculum and motor cortex activities predict motor recovery in moderate to severe stroke



Firdaus Fabrice Hannanu, Thomas A. Zeffiro, Laurent Lamalle, Olivier Heck, Antoine Thuriot, Alexandre Krainik, Marc Hommel, Olivier Detante, Assia Jaillard, ISIS-HERMES Study Group, K. Garambois, M. Barbieux-Guillot, I. Favre-Wiki, S. Grand, J.F. Le Bas, A. Moisan, M.J. Richard, F. De Fraipont, J. Gere, S. Marcel, W. Vadot, G. Rodier, D. Perennou, A. Chrispin, P. Davoine, B. Naegele, P. Antoine, I. Tropres, F. Renard

PII: S2213-1582(17)30023-2
DOI: doi: [10.1016/j.nicl.2017.01.023](https://doi.org/10.1016/j.nicl.2017.01.023)
Reference: YNICKL 933

To appear in: *NeuroImage: Clinical*

Received date: 9 November 2016
Revised date: 9 January 2017
Accepted date: 22 January 2017

Please cite this article as: Firdaus Fabrice Hannanu, Thomas A. Zeffiro, Laurent Lamalle, Olivier Heck, Antoine Thuriot, Alexandre Krainik, Marc Hommel, Olivier Detante, Assia Jaillard, ISIS-HERMES Study Group, K. Garambois, M. Barbieux-Guillot, I. Favre-Wiki, S. Grand, J.F. Le Bas, A. Moisan, M.J. Richard, F. De Fraipont, J. Gere, S. Marcel, W. Vadot, G. Rodier, D. Perennou, A. Chrispin, P. Davoine, B. Naegele, P. Antoine, I. Tropres, F. Renard, Parietal operculum and motor cortex activities predict motor recovery in moderate to severe stroke. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Ynickl(2017), doi: [10.1016/j.nicl.2017.01.023](https://doi.org/10.1016/j.nicl.2017.01.023)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Parietal operculum and motor cortex activities predict motor recovery in moderate to severe stroke

Firdaus Fabrice Hannanu^{1,2}, Thomas A. Zeffiro^{2,3}, Laurent Lamalle^{1,4-6}, Olivier Heck^{7,8}, Antoine Thuriot^{9,11}, Alexandre Krainik^{1,4-8}, Marc Hommel^{2,9,10}, Olivier Detante^{2,8,11} and Assia Jaillard^{1,2,9} *on behalf of ISIS-HERMES study group.*

- (1)Unité IRM 3T-Recherche- UMS *IRMaGe* – Centre Hospitalier Universitaire (CHU) Grenoble Alpes - France
- (2)Laboratoire MATICE - Pôle Recherche – CHU Grenoble-Alpes
- (3)Neurometrika, Potomac, MD, United States;
- (4)*IRMaGe* - Inserm US-017
- (5)*IRMaGe* - CNRS UMS-3552,
- (6)*IRMaGe* - Université Grenoble-Alpes - France
- (7)Neuroradiologie et IRM- Centre Hospitalier Universitaire Grenoble -Alpes
- (8)Grenoble Institut des Neurosciences (GIN) Inserm U836-UJF-CEA-CHU.
- (9)AGEIS, EA-UGA 7407 Université Grenoble Alpes
- (10) Clinattec - CHU Grenoble-Alpes
- (11) Unité neurovasculaire - CHU Grenoble-Alpes

Corresponding author: Assia Jaillard

Corresponding author's address: Unité IRM 3T Recherche - CHU Grenoble-Alpes - CS 10217 - 38043 Grenoble - France

Corresponding author's Tel: +33 476769306; Fax: +33 47669305

Corresponding author's e-mail address: Assia.Jaillard@ujf-grenoble.fr

(assia.jaillard@gmail.com)

Download English Version:

<https://daneshyari.com/en/article/8688731>

Download Persian Version:

<https://daneshyari.com/article/8688731>

[Daneshyari.com](https://daneshyari.com)