


# Accepted Manuscript

High incidence of deep vein thrombosis in the perioperative period of neurosurgical patients

Manabu Natsumeda, M.D. Ph.D., Takeo Uzuka, M.D. Ph.D., Jun Watanabe, M.D., Masafumi Fukuda, M.D. Ph.D., Yasuhisa Akaiwa, M.D. Ph.D., Kazuhiko Hanzawa, M.D. Ph.D., Masahiko Okada, M.D., Ph.D., Makoto Oishi, M.D. Ph.D., Yukihiro Fujii, M.D. Ph.D.



PII: S1878-8750(17)32255-6

DOI: [10.1016/j.wneu.2017.12.139](https://doi.org/10.1016/j.wneu.2017.12.139)

Reference: WNEU 7152

To appear in: *World Neurosurgery*

Received Date: 29 September 2017

Revised Date: 20 December 2017

Accepted Date: 21 December 2017

Please cite this article as: Natsumeda M, Uzuka T, Watanabe J, Fukuda M, Akaiwa Y, Hanzawa K, Okada M, Oishi M, Fujii Y, High incidence of deep vein thrombosis in the perioperative period of neurosurgical patients, *World Neurosurgery* (2018), doi: 10.1016/j.wneu.2017.12.139.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

High incidence of deep vein thrombosis in the perioperative period of neurosurgical patients

Manabu Natsumeda M.D. Ph.D.,<sup>1</sup>\* Takeo Uzuka M.D. Ph.D.,<sup>2</sup>\* Jun Watanabe M.D.,<sup>1</sup> Masafumi Fukuda M.D. Ph.D.,<sup>1,3</sup> Yasuhisa Akaiwa M.D. Ph.D.,<sup>4</sup> Kazuhiko Hanzawa M.D. PH.D.<sup>5</sup> Masahiko Okada, M.D., Ph.D.,<sup>6</sup> and Makoto Oishi M.D. Ph.D.<sup>1</sup>, Yukihiro Fujii M.D. Ph.D.,<sup>1</sup>

\* = contributed equally

<sup>1</sup>Department of Neurosurgery, Brain Research Institute, University of Niigata, Niigata, Japan

<sup>2</sup>Department of Neurosurgery, Dokkyo Medical University, Tochigi, Japan

<sup>3</sup>Department of Neurosurgery, Epilepsy Center, Nishi-Niigata Chuo National Hospital, Niigata, Japan

<sup>4</sup>Department of Neurology, Brain Research Institute, University of Niigata, Niigata, Japan

<sup>5</sup>Department of Thoracic and Cardiovascular Surgery, School of Medicine, University of Niigata, Niigata, Japan

<sup>6</sup>Department of Laboratory Medicine, School of Medicine, University of Niigata, Niigata, Japan

Key Words: venous thromboembolism; deep vein thrombosis; perioperative period; neurosurgery

Correspondence to: Yukihiro Fujii, Department of Neurosurgery, Brain Research Institute, University of Niigata, 1-757 Asahimachi, Chuo-ku, Niigata 951-8585, Japan

Tel: 025-227-0650

Fax: 025-227-0819

E-mail: yfujii@bri.niigata-u.ac.jp

Running Title: High incidence perioperative DVT in neurosurgery

Download English Version:

<https://daneshyari.com/en/article/8691853>

Download Persian Version:

<https://daneshyari.com/article/8691853>

[Daneshyari.com](https://daneshyari.com)