

**CASE REPORT**

Effect of Acupuncture on Post-implant Paresthesia

Crischina Branco Marques Sant'Anna ^{1,*},
Paulo Renato Junqueira Zuim ¹, Daniela Atili Brandini ²,
Aimée Maria Guiotti ¹, Joao Batista Vieira ³,
Karina Helga Leal Turcio ¹

¹ Department of Dental Materials and Prosthodontics, Dental School of Araçatuba, Univ Estadual Paulista—UNESP, Brazil

² Department of Integrated Clinic, Dental School of Araçatuba, Univ Estadual Paulista—UNESP, Brazil

³ Department of Pathology and Clinical Propedeutics, Dental School of Araçatuba, Univ Estadual Paulista—UNESP, Brazil

Available online ■ ■ ■

Received: Aug 22, 2016

Revised: Dec 6, 2016

Accepted: Jan 6, 2017

KEYWORDS

acupuncture therapy;
dental implants;
facial pain;
paresthesia

Abstract

Paresthesia is defined as an alteration in local sensibility, associated with numbness, tingling, or unpleasant sensations caused by nerve lesions or irritation. It can be temporary or permanent. The treatment protocol for facial paresthesia is primarily based on the use of drugs and implant removal, which may not be completely effective or may require other risk exposure when there is no spontaneous regression. However, other therapeutic modalities such as acupuncture can be used. The aim of this study is to report a case of a patient with paresthesia of the inferior alveolar nerve and pain caused by an implant surgery performed 2 years earlier. The patient received acupuncture treatment during 4 months of weekly sessions. Six points were used: Large Intestine (LI4), Large Intestine (LI11), Stomach (ST36), Liver (LR3), Extra Head and Neck (E-HN-18), and Stomach (ST5). The visual analog scale was used before and after each session for the analysis of paresthesia and pain, together with assessment of the paresthesia by delimitation of the desensitized region of the skin and presented discomfort. Pain remission and reduction in the size of the paresthesia area occurred after four sessions.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

* Corresponding author. Aguapei, 3300, Araçatuba, São Paulo, CEP 16025-455, Brazil.

E-mail: crischina.s@terra.com.br (C.B.M. Sant'Anna).

pISSN 2005-2901 eISSN 2093-8152

<http://dx.doi.org/10.1016/j.jams.2017.01.004>

Copyright © 2017, Medical Association of Pharmacopuncture Institute.

1. Introduction

Surgical procedures have increased in dentistry as has the risk of consequences, such as paresthesia (3.6–11.9%) [1], with transitional (23%) or persistent complications (13%) [2,3], with 64.4% of these in the inferior alveolar nerve [4]. Paresthesia is a sensibility disorder with strange or unpleasant sensations. It can produce a partial loss of sensitivity or spontaneous abnormal sensations such as tingling. It is caused by mechanical factors, such as direct or indirect trauma, and pathological, physical, chemical, and microbiological alterations. It is a traumatic, painful, peripheral postimplant neuropathy, which may be associated with hypoesthesia, hyperalgesia, dysesthesia, allodynia, and pain. The amount of alteration is proportional to the fiber type, amount of tissue loss, and intensity of injury suffered [5].

Imaging tests are important to prevent accidents [6]. Medications to reduce postoperative edema and inflammation are also preconized. These include nonsteroidal anti-inflammatory drugs, injectable steroids, and physical resources, such as heat, transcutaneous electrical nerve stimulation, lasers, and acupuncture [7–9].

According to Traditional Chinese Medicine (TCM), the organism works correctly when there is a dynamic balance between two fundamental aspects: Qi (energy) and matter (organic part) [10,11]. The energy is absorbed and distributed throughout the organism through energy channels, and the material portion is distributed through the circulatory system [10]. All innate factors, unruly feeding, and suppressed or intense emotion can lead to disharmony of the systems, provoking illness [10–12]. Surgical trauma causes blockages that cause pain [10,11], tingling, and heavy (numbness) sensations, which are characteristics of paresthesia [13]. By TCM theory, when inserting needles in specific points, analgesic mechanisms are unleashed [14] through the release of endogenous substances, improving the tissue healing response, nerve conduction [15], and local blood flow [16]. Based on the TCM perspective, paresthesia results from the stagnation and blocking of the Qi (energy or flow of nerve signals) [12].

The aim of this study is to report a case of a patient with paresthesia of the inferior alveolar nerve caused by an implant surgery performed 2 years earlier. The patient received acupuncture treatment for the condition.

2. Case presentation

A 56-year-old female patient came to the Temporomandibular Disorders Diagnosis and Treatment Center (TMD) at the Aracatuba Dental School, Araçatuba, São Paulo, Brazil, reporting a strange sensation in the lower lip and in the delimited region of the mental foramen and mucosa hemiarch [from the median line up to the left mandibular second molar (Tooth 36)]. She also reported acute pain in the mucosa at a specific point within the paresthesia area (region of Tooth 35). The symptoms appeared after implant surgeries for Teeth 36 and 37 (Fig. 1).

The patient reported pain during suturing and subsequent removal. While yawning 3 days after the surgery, she felt an acute sensation in the region that developed into a



Aspect of the dysesthesia after acupuncture



Aspect of the dysesthesia before acupuncture

Figure 1 Dysesthesia before and after the acupuncture treatment in the green area.

burning sensation which occurred sporadically while brushing teeth, talking, or sometimes spontaneously. Over time, this area developed a painful, unpleasant sensation described as numbness (paresthesia).

The predominance of the earth element, which includes discomfort with humidity, preference for sweet foods, presence of anxiety and preoccupation, yellowing of the skin, and darkening around the eyes, was observed in the patient. Additionally, a normal colored, tapered tongue without coating, which had a slight deviation to the right, a red tip, and indentations on the side, was observed.

The patient did not use medication for the symptoms. The study was conducted in accordance with ethical criteria and the Declaration of Helsinki.

Weekly acupuncture sessions were held for 4 months. Six points were used: Large Intestine (LI4), Large Intestine (LI11), Stomach (ST36), Liver (LR3), Extra Head and Neck (EHN-18), and Stomach (ST5). These points were selected according to the characteristics of the patient and the clinical features observed (paresthesia and muscle pain—TMD) [10]. The visual analog scale was used before and after each session for the analysis of paresthesia and pain, together with the assessment of the paresthesia by delimitation of the desensitized region of the skin and presented discomfort.

After the first session, a decrease in the intensity of pain was observed. After three sessions, acute pain became less frequent, disappearing after the fourth session.

Download English Version:

<https://daneshyari.com/en/article/8692600>

Download Persian Version:

<https://daneshyari.com/article/8692600>

[Daneshyari.com](https://daneshyari.com)