




Contents lists available at ScienceDirect

Chinese Journal of Traumatology

journal homepage: <http://www.elsevier.com/locate/CJTEE>

Original Article

Early scaphoid fractures are better diagnosed with ultrasonography than X-rays: A prospective study over 114 patients

Ravikant Jain ^a, Nikhil Jain ^{b,*}, Tanveer Sheikh ^a, Charanjeet Yadav ^c^a Department of Orthopedics, SAMC and PGI, Indore, MP, India^b Department of Orthopedics, MLB Medical College, Jhansi, UP, India^c Department of Radio Diagnosis, SAMC and PGI, Indore, MP, India

ARTICLE INFO

Article history:

Received 14 August 2017

Received in revised form

10 November 2017

Accepted 14 November 2017

Available online xxx

Keywords:

Scaphoid fractures

Ultrasonography

High frequency

ABSTRACT

Purpose: Wrist has a complex anatomy and undergoes complex injuries. Scaphoid fracture is one of such injuries. It is the most common fracture in carpal bone. Most of the scaphoid fractures are missed on initial X-rays. Magnetic resonance imaging (MRI) is considered as a gold standard for diagnosing scaphoid fractures. Ultrasonography (USG) is emerging as a good alternative to make an early diagnosis of scaphoid fractures. Our aim is to throw light upon the role of USG in detection of scaphoid fractures. **Methods:** The study was centered upon 114 patients in the age range 10–65 years, with traumatic wrist injury and were clinically suspected to have scaphoid fractures. Patient with non-traumatic history, bilateral wrist injury and late presentation were excluded. X-rays, USG using high frequency probe and MRI were done for all patients. MRI was considered to be the gold standard test. Patients were followed up at 6 weeks.

Results: Of the 114 patients, X-ray could diagnose scaphoid fractures in 48 patients, 30 of which were confirmed by MRI. USG results were positive in 74 patients, of which MRI was positive in 67 patients. The accuracy of scaphoid fracture detection with USG was 98.04% in comparison to X-ray (20.58%), which was statistically significant.

Conclusion: USG provides a more accurate and reliable method of making an early diagnosis of scaphoid fracture than X-rays. It is non-invasive, non-expensive and allows better visualisation of cortical disruption.

© 2018 Daping Hospital and the Research Institute of Surgery of the Third Military Medical University.

Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The hand and wrist constitute very important functional parts of the body used in daily life activities. So these are very much vulnerable to traumatic injuries. 6.6%–28.6% of all injuries and 28.0% of all musculoskeletal injuries comprise injury to wrist and hand.^{1,2} Wrist injuries are a bit commoner than hand injuries. Schoffl et al³ reported that these injuries accounted for 14.0%–30.0% of all patients being treated in emergency room. Though these injuries never threaten life, early diagnosis is always necessary, so that the appropriate treatment can be started as early as possible. Accurate diagnosis can be made with proper clinical examination and should be confirmed with imaging before any treatment is advised.

Radiographs remain the investigation of choice for arriving at diagnosis when injury occurs. Specialised views are required to visualise all aspects of wrists and hand. Above all, clinical correlation is equally significant. If the findings on radiographs and clinical examination are not correlating, one should prefer other imaging modalities. Ultrasonography (USG), magnetic resonance imaging (MRI), and computed tomography (CT) are other such imaging techniques, which provide accurate diagnosis that are often missed on X-rays. Many studies consider that with MRI, occult scaphoid fractures could be diagnosed earlier.^{4–6} MRI has a high sensitivity and specificity for scaphoid fractures not evident on X-rays.^{7,8} Some consider MRI as the gold standard investigation for clinically suspected scaphoid fracture, and it allows diagnosis of occult bony and soft-tissue injuries.⁹ High frequency USG has also been recommended to diagnose occult scaphoid fractures.¹⁰ The criteria used for USG detection of scaphoid fractures were cortical disruption and wrist effusion.¹¹

* Corresponding author.

E-mail address: nikrockin23@gmail.com (N. Jain).

Peer review under responsibility of Daping Hospital and the Research Institute of Surgery of the Third Military Medical University.

<https://doi.org/10.1016/j.cjtee.2017.09.004>1008-1275/© 2018 Daping Hospital and the Research Institute of Surgery of the Third Military Medical University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The objective of our study is evaluation of traumatic wrist injuries using ultrasonography, and making an early diagnosis of scaphoid fractures. Scaphoid is the most frequently fractured carpal bone.¹² The most common mechanism of injury is a fall onto an outstretched hand. Wrist pain, swelling and tenderness over anatomical snuffbox on clinical examination raise the suspicion of scaphoid fracture. Most of the scaphoid fractures are occult on presentation and often missed on initial radiographs. USG is economic and a non-invasive imaging modality, which uses higher frequency sound waves that are normally not audible to human beings.¹³ These higher frequencies provide better resolution and improved spatial detail. We have evaluated 114 patients with traumatic wrist and hand injuries in our study and they were subjected to digital X-ray, USG, and MRI. MRI has been considered as gold standard in our study.

Materials and methods

A total of 146 patients in the age group of 10–65 years with traumatic wrist injuries were evaluated. Of these, 114 agreed to participate in the study. Written and informed consents were taken. The patients presented with complaints of pain, swelling or restricted movements of wrist joint or tenderness in anatomical snuffbox in the outdoor patient department of author's institute and had a clinical suspicion of scaphoid fracture. Patients with bilateral involvement, non-traumatic history and presenting after a week from trauma were excluded from the study. Detailed history was taken and bilateral wrist joint examination was done. Provocative tests included Scaphoid shift test (pain on volar-dorsal shifting of the scaphoid) and Watson shift test (pain on moving wrist from ulnar to radial deviation). Digital radiographs of involved wrist joint were then obtained. Standard radiographic views, neutral anteroposterior and lateral along with some special views for scaphoid, pronated oblique and posteroanterior in ulnar deviation of wrist were preferred. Advantages of views in two perpendicular planes are easy identification of bony landmarks and assessment of adjacent soft tissue structures.

Patients were then subjected to ultrasonography (USG) of the affected wrist. All USGs were reviewed by single radiologist to avoid intra-observer variability. The healthy side was used as control. USG was used along with extended field of view (EFOV) and colour Doppler technology whenever required. Use of such technologies allows clear visualisation of bones, joints, muscles, tendons, and even slightest inflammatory conditions. A linear array transducer of frequency 5–12 MHz (Million Hertz) was used in our study.

Patients were made to sit on a stool/couch facing the USG machine and asked to expose both the wrist joints up to the mid forearm simultaneously. Hand was placed appropriately for imaging specific areas of the wrist. Longitudinal, transverse, and EFOV images were obtained. The radiologist was asked to assess the cortical continuity of scaphoid and detect the presence of radio-carpal effusion. Signs suggestive of scaphoid fracture on USG were focal defect due to cortical disruption of scaphoid or hypoechoic fluid in the radio carpal space due to haemarthrosis. These signs were reported to be associated with scaphoid fractures.^{11,14,15} Fig. 1 shows scaphoid fractures detected with USG in patients with normal X-rays.

All patients were then subjected to dedicated wrist MRI, which was considered to be the gold standard.

Results

The average age of patients was 32 years (range, 10–65 years), with 65% patients between 20 and 40 year, 15% below 10 years, and rest 20% above 40 years. Male: female ratio was 2:1.75% of the patients were from the middle socioeconomic class, and remaining 25% were from the lower class. The chief complaints were pain, swelling and restricted wrist movements. Most of the patient presented with more than one complaint. 2% of the patient had deformity.

Digital X-rays, USG and MRI of the involved wrist joint were ordered for all patients upon initial presentation. Of the 114, 48 patients were diagnosed to have scaphoid fracture on X-ray




Fig. 1. A: Initial X-ray with normal findings. B: USG showing cortical disruption as pointed by arrow head. C: USG showing decreased width of fractured scaphoid due to impaction as compared to normal side (blue line-normal side, green line-affected side).

Download English Version:

<https://daneshyari.com/en/article/8694864>

Download Persian Version:

<https://daneshyari.com/article/8694864>

[Daneshyari.com](https://daneshyari.com)