



Contents lists available at ScienceDirect

Chinese Journal of Traumatology

journal homepage: <http://www.elsevier.com/locate/CJTEE>

Case report

Innervated full thickness grafts in distal finger amputations

Anouk J.M. Cornelissen^{*}, Martijn A. van Onna, Xavier H.A. Keuter, René RWJ. van der Hulst

Maastricht University Medical Center, Department of Plastic and Reconstructive Surgery, P. Debyelaan 25, Maastricht 6229 HX, The Netherlands

ARTICLE INFO

Article history:

Received 12 December 2016

Received in revised form

3 May 2017

Accepted 13 June 2017

Available online xxx

Keywords:

Finger joint

Amputation

Surgical flaps

Innervation

Neuroma

ABSTRACT

Innervated full thickness graft will be presented as an option to reconstruct a fingertip defect which might result in better sensibility than standard reconstruction using a full thickness graft without innervation. Also, anastomosing the nerve stumps can decrease the chance of developing neuroma.

© 2017 Daping Hospital and the Research Institute of Surgery of the Third Military Medical University.

Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The usage of an innervated full thickness graft was first described by Lister in 1978 for coverage of a heel defect.¹ However, as far as we know an innervated full thickness graft for the coverage of fingertip amputations has not been used before. In literature only complex methods for a sensible reconstruction of a fingertip are described, e.g. a partial toe transfer, a wraparound flap or a neurovascular pedicle V-Y advancement flap.^{2,3} We would like to present two cases in which a more simple technique was used to reconstruct a fingertip with innervation. An innervated full thickness graft for a fingertip was used.

Case report

The first case describes a fifteen-year-old boy who got injured when he was using a circular saw, working on a crafts project. He presented at the emergency room with extensive trauma of the left thumb and index finger. The thumb showed a dorsal laceration with a lesion of the extensor pollicis longus tendon and a fracture of the proximal phalanx. On the index finger a ragged amputation at the distal interphalangeal joint with partial deglovement of the

radial side was found, the flexor digitorum profundus was not functional. X-ray examination revealed an inverted T-shaped intra-articular fracture of the proximal phalanx of the thumb and an amputation of the distal phalanx of the index finger.

During the operation the thumb fracture was fixed using a T-shaped osteosynthesis plate using screws, the extensor tendon was repaired using a two-strand modified Kessler and the wound was closed using transcutaneous sutures.

Due to the distal level and raggedness of the amputation of the index finger a re-implantation was not feasible. The partial deglovement was covered using the skin of the amputated part as a full thickness graft of approximately 3 cm.² During the process of defatting the graft the radial digital nerve of the index finger (RDN2) was identified and freed. The nerve was freed using micro-instruments and glasses with surgical loops. The proximal part of the RDN2 was also identified and freed using the same instruments. An anastomosis of the RDN2 was made using 9-0 sutures through the epineurium. The transplant was sutured into place using transcutaneous sutures and a cast was provided (Figs. 1 and 2).

The second case describes a 66-year-old male with an amputation of the third digit at the level of the midphalanx and of the fourth digit at the level of the DIP joint after removing loose grass from a running lawnmower. The amputated part of the third digit was missing and probably shredded as an additional search at the site of injury did not retrieve the fragment. The amputated part of the fourth digit was partially shredded. In both digits good function of the flexor digitorum superficialis was found but

^{*} Corresponding author.

E-mail address: c.anouk@hotmail.com (A.J.M. Cornelissen).

Peer review under responsibility of Daping Hospital and the Research Institute of Surgery of the Third Military Medical University.

<https://doi.org/10.1016/j.cjtee.2017.04.008>

1008-1275/© 2017 Daping Hospital and the Research Institute of Surgery of the Third Military Medical University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Fig. 1. Pre- and postoperative pictures of the injured and reconstructed hand of the first case.



Fig. 2. Postoperative picture at 18 months of the first case. This patient tattooed a smiley a pair of scissors and an arrow to the injured finger 12 months after the operation as closure of the treatment.

nerve of the fourth digit (RDN4) was identified in the amputated part and was anastomosed to the proximal part of the radial digital nerve (RDN3) with 9-0 sutures through the epineurium. Surgical technique was comparable to the technique used in the first case. The transplant was sutured into place using transcutaneous sutures (Figs. 3 and 4).



Fig. 4. Postoperative picture at 12 months of the second case.

disfunctioning of the flexor digitorum profundus was noticed. In addition, there was a large soft tissue defect of the third digit. X-ray examination showed a fracture of the amputated distal phalanx and a small part of the medial phalanx. The third digit was amputated in the middle of the mid phalanx with a small fracture of the distal part of the remaining bone. During the operation both digits were slightly shortened and the fracture was fixated. The fourth digit was closed primarily. The third digit could not be closed primarily because of a large soft tissue defect. The shredded amputate of the fourth digit was used as a full thickness graft to cover the defect which measured around 2.5 cm². The radial digital



Fig. 3. Pre- and postoperative pictures of the injured and reconstructed hand of the second case.

Download English Version:

<https://daneshyari.com/en/article/8694935>

Download Persian Version:

<https://daneshyari.com/article/8694935>

[Daneshyari.com](https://daneshyari.com)