

Accepted Manuscript

Axial four-dimensional computed tomographic images to analyze crosswise differences in protrusive condylar movement in patients who underwent mandibulectomy and free flap reconstruction

Masaya Akashi, DDS, PhD, Assistant Professor, Toshinori Sekitani, radiologic technologist Clinical fellow, Yumi Ohtsuki, DDS, Clinical fellow, Yasumasa Kakei, DDS, PhD, Assistant Professor, Junya Kusumoto, DDS, Clinical fellow, Takumi Hasegawa, DDS, PhD, Assistant Professor, Michinori Maeda, DDS, Clinical fellow, Noriyuki Negi, radiologic technologist Clinical fellow, Kazunobu Hashikawa, MD, PhD, Associate Professor, Yasuyuki Shibuya, DDS, PhD, Professor and Chairman, Satoru Takahashi, MD, PhD, Director and Professor, Takahide Komori, DDS, PhD, Professor and Chairman

PII: S1010-5182(17)30294-9

DOI: [10.1016/j.jcms.2017.08.025](https://doi.org/10.1016/j.jcms.2017.08.025)

Reference: YJCMS 2770

To appear in: *Journal of Cranio-Maxillo-Facial Surgery*

Received Date: 6 February 2017

Revised Date: 27 July 2017

Accepted Date: 23 August 2017

Please cite this article as: Akashi M, Sekitani T, Ohtsuki Y, Kakei Y, Kusumoto J, Hasegawa T, M, Negi N, Hashikawa K, Shibuya Y, Takahashi S, Komori T, Axial four-dimensional computed tomographic images to analyze crosswise differences in protrusive condylar movement in patients who underwent mandibulectomy and free flap reconstruction, *Journal of Cranio-Maxillofacial Surgery* (2017), doi: 10.1016/j.jcms.2017.08.025.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Axial four-dimensional computed tomographic images to analyze crosswise differences in
protrusive condylar movement in patients who underwent mandibulectomy and free flap
reconstruction**

Masaya Akashi, DDS, PhD,^{a†*} Toshinori Sekitani, RT,^{b‡} Yumi Ohtsuki, DDS,^{a‡} Yasumasa Kakei, DDS,
PhD,^{a‡} Junya Kusumoto, DDS,^{a‡} Takumi Hasegawa, DDS, PhD,^{a‡} Michinori Maeda, DDS,^{c‡} Noriyuki
Negi, RT,^{b‡} Kazunobu Hashikawa, MD, PhD,^{d‡} Yasuyuki Shibuya, DDS, PhD,^{e‡} Satoru Takahashi, MD,
PhD,^{b§} Takahide Komori, DDS, PhD,^{a¶}

† Assistant Professor, ‡ Clinical fellow, ¶ Associate Professor, § Director and Professor, ¶ Professor and

Chairman

RT: radiologic technologist.

^aDepartment of Oral and Maxillofacial Surgery, Kobe University Graduate School of Medicine (Chief: Prof.

Komori Takahide, DDS, PhD), Kobe, Japan

^bDepartment of Radiology, Kobe University Hospital (Chief: Prof. Satoru Takahashi, MD, PhD), Kobe, Japan

^cDepartment of Oral and Maxillofacial Surgery, Nagoya City University Graduate School of Medical

Sciences (Chief: Prof. Yasuyuki Shibuya, DDS, PhD), Nagoya, Japan

Download English Version:

<https://daneshyari.com/en/article/8699040>

Download Persian Version:

<https://daneshyari.com/article/8699040>

[Daneshyari.com](https://daneshyari.com)