

Contents lists available at [ScienceDirect](#)

Journal of Dental Sciences

journal homepage: www.e-jds.com

Original Article

Quantitative analysis of Miller mobility index for the diagnosis of moderate to severe periodontitis - A cross-sectional study

Chung-Ping Wu ^a, Yu-Kang Tu ^b, Sao-Lun Lu ^a, Jui-Hung Chang ^a, Hsein-Kun Lu ^{a, c, *}^a Periodontal Department, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan^b Institute of Epidemiology and Preventive Medicine, College of Public Health, National Taiwan University, Taipei, Taiwan^c Clinical Periodontics of Dental Department, Taipei Medical University Hospital, Taipei, Taiwan

ARTICLE INFO

Article history:

Received 7 September 2017

Received in revised form

16 November 2017

Accepted 17 November 2017

Available online xxx

Keywords:

Miller's mobility index

Negative predictive value

Nonsurgical periodontal therapy

Positive predictive value

Receiver operating characteristic curve

Tooth mobility

ABSTRACT

Background/purpose: We assessed the mobility of single-root teeth by using Miller's mobility index (MMI) and to analyze the validity of MMI for the diagnosis of periodontitis.

Materials and methods: A total of 30 patients were included and the Spearman correlation coefficient was used to assess the correlation between MMI, clinical attachment level (CAL), and probing depth (PD). The validity of MMI for the diagnosis of the severity of periodontitis was evaluated using the receiver operating characteristic (ROC) curve, area under curve (AUC) value, positive predictive value (PPV).

Results: Strong correlations were observed between MMI and CAL ($r = 0.92$) and between MMI and PD ($r = 0.76$). When the CAL = 3–4 mm and CAL ≥ 5 mm groups were pooled together, the AUC value was 0.81. The AUC was 0.86 for diagnosis with MMI in the CAL ≥ 5 mm group. A PPV of 100% was achieved for all grades when MMI > 1 .

When the teeth with PD ≥ 5 to < 7 mm and PD ≥ 7 mm groups were pooled together, the AUC value for MMI was 0.80. The PPV was 98.8%, 99%, and 100% for MMI Grade 1, Grade 2, and Grade 3, respectively. When PD ≥ 7 mm was defined as severe periodontitis, the AUC value for MMI was 0.72.

Conclusion: MMI may provide valuable information for the diagnosis of moderate and severe periodontitis when CAL is not obtainable during routine practice.

© 2018 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Periodontitis is an inflammatory reaction caused by bacteria. It can lead to the destruction of the gingiva and connective tissues, and, in particular, alveolar bone loss. If left untreated, it ultimately leads to tooth loss. Increased tooth mobility is a common symptom of severe periodontal disease and is one of the main reasons patients seek consultations at periodontal clinics. A study conducted by Eke on the prevalence of periodontitis in adults aged 30 years and older in the United States demonstrated that in 2009–2012, 46% of adults in the United States, or approximately 64.7 million adults, had periodontitis, with 8.9% of cases being severe.¹ Between 2003 and 2005, Lai conducted a study on residents of Keelung, Taiwan who were aged 35–44 years to assess the prevalence of

periodontal disease. The community periodontal index and loss of clinical attachment were used as parameters, and the prevalence of periodontal disease was determined to be 94.8%.²

Despite increasing understanding of the reasons for and mechanisms of periodontal disease in recent years, the methods used by dentists to diagnose periodontal disease have not changed. Diagnoses are still largely based on dental X-rays and clinical periodontal parameters such as probing depth (PD), clinical attachment level (CAL), bleeding on probing (BOP), and tooth mobility. Miller's mobility index (MMI) is the most widely accepted method for routine clinical examinations of tooth mobility.³ The tooth is held between the metallic handles of two instruments and moved in the buccolingual or buccopalatal direction, and the moved distance is visually estimated by the person conducting the examination. Mobility is then classified into Grades 0–3. Although special devices such as periodontometers can be used to obtain quantitative measurements of tooth mobility, the operation of such devices is complicated and time consuming.⁴

In recent years, receiver operating characteristic (ROC) curves

* Corresponding author. author. Periodontal Department, College of Oral Medicine, Taipei Medical University; 250 Wu Xing Street, Taipei, Taiwan.

E-mail address: Jackson@tmu.edu.tw (H.-K. Lu).

<https://doi.org/10.1016/j.jds.2017.11.001>

1991-7902/© 2018 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

have been widely used to diagnose diseases.⁵ The true positive rate, which is indicated as sensitivity, is plotted on the y-axis of the ROC curve, and the false positive rate, which is indicated as 1-specificity, is plotted on the x-axis. A 45° diagonal line is used as a reference line for interpreting the ROC curve. The closer the ROC curve is to the upper left corner, the higher the sensitivity of the diagnostic tool toward the disease and the lower the false positive rate, indicating that this diagnostic tool exhibits more efficient discrimination than others.

The purpose of this study was to assess quantitative changes in the mobility of single-rooted anterior tooth by using MMI and to analyze the correlation between mobility and various clinical periodontal parameters. The ROC, area under curve (AUC), sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) of MMI for the clinical diagnosis of periodontal disease were assessed.

Materials and methods

Patients with no systemic diseases who were visiting the Periodontal Clinic of the Department of Dentistry at Taipei Medical University Hospital for the first time and who had not received previous treatment for periodontal disease at other hospitals or clinics were recruited. The patients were diagnosed as having full-mouth chronic periodontitis, with at least six anterior teeth with mean PD \geq 5 mm in combination with BOP (+), and they received complete nonsurgical periodontal therapy. Patients with a history of AIDS, unstable cardiovascular disease, and uncontrolled diabetes mellitus (HbA1c > 8.0%) were excluded from the study. In addition, pregnant women and people with mental or intellectual disabilities, parafunction of chewing habit, severe malocclusion, and those who received clinical treatment and antibiotic therapy for periodontal disease in the past 6 months were excluded.

This study was approved by the Taipei Medical University Joint Institutional Review Board, and a certificate of approval (No. N201512036) has been issued. All participants were informed of the significance and purpose of the study and signed a consent form.

A total of 30 patients (13 men and 17 women) who had received a diagnosis of moderate to severe periodontitis from April 20, 2016 onwards December 20, 2016 were selected. The clinical periodontal parameters included MMI, CAL, PD, BOP, and plaque index (PI). Six sites for each anterior tooth (tooth #13–23 and #33–43) without obvious fremitus in the position of centric occlusion were measured.

The Williams true pressure-sensitive probe (Pro-Dentec, Arkansas, USA) was used to measure PD and CAL. All examinations were performed by a highly trained dentist (Wu CP), and intra-examiner kappa values for PD and CAL measurements were evaluated before the investigation.

According to the American Academy of Periodontology (AAP) 2015 Task Force Report, BOP (+), PD, and CAL measurements are necessary for the classification of mild, moderate, and severe periodontitis. CAL = 3–4 mm or PD \geq 5 to <7 mm were defined as moderate periodontitis and CAL \geq 5 mm or PD \geq 7 mm were defined as severe periodontitis.⁶

Griner et al. revealed that statistical validation of a potentially useful diagnostic test involved the use of a 2 × 2 decision matrix.⁷ Accordingly, the use of ROC, AUC, sensitivity, specificity, PPV, and NPV of MMI as diagnostic tools for periodontal disease were assessed. Spearman correlation coefficient was used to assess the correlation between MMI, CAL and PD. Statistical analysis was performed using SPSS version 19.0 (SPSS Inc., Chicago, USA). The level of significance was set at $p < .05$.

Table 1
Basic data in gender.

	Total	Male	Female
Person number	30	13	17
Tooth number	338	152	186
Mean age (years)	60.45	59.68	61.03
Age range (years)	46–70	46–70	46–70

Table 2
Correlation between MMI and other parameters in gender.

	30 Subjects		13 Male		17 Female	
	correlation	r	correlation	r	correlation	r
MMI/CAL	high	0.92	high	0.94	high	0.91
MMI/PD	high	0.76	high	0.82	high	0.71
MMI/PI	medium	0.31	low	0.20	medium	0.36
MMI/BOP	medium	0.35	medium	0.39	medium	0.33

MMI = Miller's mobility index.
CAL = clinical attachment level.
PD = probing depth.
BOP = bleeding on probing.

Results

The intra-examiner kappa values for PD and CAL were 0.93 and 0.88, respectively. Table 1 presents the baseline characteristics of all participants. A total of 30 patients (13 men and 17 women) with a mean age of 60.45 years and 338 teeth (152 belonging to men, 186 belonging to women) were included in this study.

The correlations between MMI and various clinical periodontal parameters for different gender groups are presented in Table 2. A strong correlation was observed between MMI and CAL ($r = 0.92$) and between MMI and PD ($r = 0.76$).

Table 3 presents the numbers and percentages of teeth classified as having mild, moderate, and severe periodontitis among the gender groups, on the basis of various definitions of periodontitis. According to the definition of severe periodontitis as CAL \geq 5 mm, of 338 teeth, 79.29% were classified as having severe periodontitis. According to the definition of severe periodontitis as PD \geq 7 mm, only 12.42% of teeth were classified as having severe periodontitis.

When we pooled the teeth with CAL = 3–4 mm and CAL \geq 5 mm as the moderate to severe group, the ROC curve for MMI grades was obtained as shown in Fig. 1a. The curve was situated in the upper left region of the diagonal line. The AUC value was 0.81. The true positive rate was higher than the false positive rate, which indicates that MMI has an excellent discriminating power in the diagnosis of moderate to severe periodontitis. The optimal cutoff point was MMI

Table 3
Teeth numbers and percentage of various severities on. Different definition of AAP 2015 Task Force Report in gender.

	Total		Male		Female	
	numbers	%	Numbers	%	Numbers	%
Mild P.						
CAL 1–2 mm	10	2.95%	2	1.3%	8	4.3%
PD > 3 to <5 mm	36	10.65%	15	9.87%	21	11.29%
Moderate P.						
CAL 3–4 mm	60	17.75%	29	19%	31	16.67%
PD \geq 5 to <7 mm	207	61.24%	83	54.6%	124	66.67%
Severe P.						
CAL \geq 5 mm	268	79.29%	121	79.61%	39	79.03%
PD \geq 7 mm	42	12.42%	30	19.74%	12	6.45%

Mild P = mild periodontitis.
Moderate P = moderate periodontitis.
Severe P = severe periodontitis.

Download English Version:

<https://daneshyari.com/en/article/8699109>

Download Persian Version:

<https://daneshyari.com/article/8699109>

[Daneshyari.com](https://daneshyari.com)