

3. Hoorfnaegle JH, Doo E, Liang TJ, Fleischer R, Lok AS. Management of hepatitis B: summary of a clinical research workshop. *Hepatology*. 2007;45:1056–75.
4. Seto WK. Hepatitis B virus reactivation during immunosuppressive therapy: appropriate risk stratification. *World J Hepatol*. 2015;7:825–30.
5. Cantini F, Boccia S, Goletti D, Iannone F, Leoncini E, Panic N, et al. HBV reactivation in patients treated with antitumor necrosis factor- α (TNF- α) agents for rheumatic and dermatologic conditions: a systematic review and meta-analysis. *Int J Rheumatol*. 2014;2014:926836.
6. Ott JJ, Stevens GA, Groeger J, Wiersma ST. Global epidemiology of hepatitis B virus infection: new estimates of age-specific HBsAg seroprevalence and endemicity. *Vaccine*. 2012;30:2212–9.
7. Vassilopoulos D, Apostolopoulou A, Hadziyannis E, Papathodoridis GV, Manolakopoulos S, Koskinas J, et al. Long-term safety of anti-TNF treatment in patients with rheumatic diseases and chronic or resolved hepatitis B virus infection. *Ann Rheum Dis*. 2010;69:1352–5.
8. Charpin C, Guis S, Colson P, Borentain P, Mattei JP, Alcaraz P, et al. Safety of TNF-blocking agents in rheumatic patients with serology suggesting past hepatitis B state: results from a cohort of 21 patients. *Arthritis Res Ther*. 2009;11:179.

Pustular Secondary Cutaneous Aspergillosis in an Immunosuppressed Patient[☆]



Aspergillosis cutânea secundária pustulosa em paciente imunossuprimido

To the Editor:

The growing use of immunosuppressive agents in procedures such as kidney transplantation and in intensive polychemotherapy regimens for different types of cancer has led to a notable increase in opportunistic fungal infections.^{1,2} *Aspergillus* species are ubiquitous, opportunistic, filamentous fungi often found in soil, decaying organic matter, and even in food remains.² They tend to multiply in environments with high levels of dust dispersal and are particularly common in hospitals during building or maintenance work.² Care should therefore be taken to protect immunocompromised patients or patients with a greater risk of infection from exposure to building work or damp environments. *Aspergillus* species can cause serious primary or secondary skin infections.³ We present a case of pustular cutaneous aspergillosis.

A 56-year-old man with type IgA multiple myeloma was evaluated for painless skin nodules measuring over 1 cm and a large blister of recent onset on his left elbow. The patient had stage IIIA disease and had been under follow-up for 4 years. He had received several treatments, includ-

9. Lan JL, Chen YM, Hsieh TY, Chen YH, Hsieh CW, Chen DY, et al. Kinetics of viral loads and risk of hepatitis B virus reactivation in hepatitis B core antibody-positive rheumatoid arthritis patients undergoing anti-tumour necrosis factor alpha therapy. *Ann Rheum Dis*. 2011;70:1719–25.
10. European Association For The Study Of The Liver. EASL clinical practice guidelines: management of chronic hepatitis B virus infection. *J Hepatol*. 2012;57:167–85.

R. Pereira,^a I. Raposo,^b F. Nery,^{a,c} T. Torres^{a,b,*}

^a Instituto de Ciências Biomédicas Abel Salazar, University of Porto, Portugal

^b Department of Dermatology, Centro Hospitalar do Porto, Portugal

^c Hepato-Pancreatic Unit, Centro Hospitalar do Porto, Porto, Portugal

* Corresponding author.

E-mail address: torres.tiago@outlook.com (T. Torres).

1578-2190/

© 2017 Elsevier España, S.L.U. and AEDV. All rights reserved.

ing 4 cycles of chemotherapy with bortezomib 1.3 mg/m² every 4 days, 4 cycles separated by a week of cyclophosphamide 500 mg once a day for 3 days, and dexamethasone 40 mg every 2 days for 12 days. He had also received radiation therapy and undergone hematopoietic stem cell transplantation. Following a relapse in 2015, it was decided to attempt mini-allogeneic transplantation with reduced-intensity Flume-ATG conditioning (70 mg/m² melphalan, fludarabine 30 mg/m²/d, bortezomib 1.3 mg/m², and anti-thymocyte globulin 2 mg/m²) and an increase in melphalan infusion dose to 150 mg/m².

Forty days after the transplantation, the patient was evaluated by a dermatologist as he suddenly developed painless erythematous subcutaneous nodules measuring approximately 3 cm on the anterior surface of both thighs and on the left abdominal flank (Fig. 1A). One of the lesions on the lateral surface of his left elbow was a tense 1.5-cm blister containing blood-stained pus with a fluid level (Fig. 1B). The patient reported no other symptoms. The onset of these lesions coincided with a progressive increase in serum galactomannan levels, which rose from previously undetectable levels to a level of 0.9.

In view of the patient's condition and the general clinical picture, skin biopsy samples were taken for histology and microbial culture. Histologic examination of the elbow lesion showed a purulent subepidermal blister and an underlying infiltrate composed of abundant polymorphonuclear leukocytes that caused notable tissue destruction, with weakened structures, collagen bundles with an unstructured appearance, and effacement of adnexal structures (Fig. 2A). Higher magnification and periodic acid-Schiff (PAS) staining showed septate linear structures with dichotomous acute-angle (45°) branching throughout the dermis and extending into the more superficial areas of the subcutaneous tissue. These structures had an approximate diameter of 3 μ m and a length of up to 80 μ m in some sections and were consistent with hyalohyphomycosis (Fig. 2B). Culture

[☆] Please cite this article as: Fonda-Pascual P, Fernández-González P, Moreno-Arrones OM, Miguel-Gómez L. Aspergillosis cutânea secundária pustulosa em paciente imunossuprimido. *Actas Dermosifiliogr*. 2018;109:287–290.

in Sabouraud agar produced *Aspergillus flavus*, which was sensitive to voriconazole and echinocandins in the ETEST (Biomérieux).

Intensive antifungal treatment was initiated with voriconazole (loading dose of 400 mg followed by a maintenance dose of 200 mg/12 h) and intravenous anidulafungin (loading dose of 200 mg followed by 100 mg/24 h). Three days later, the patient developed right hemiparesis. In the staging study, the chest computed tomography (CT) scan showed previously undetected cavitated lesions in the right upper lobe of the lung (Fig. 3A). The brain CT scan showed 2 nonvascular frontal lesions consistent with a stroke secondary to infection (Fig. 3B). We decided to escalate the treatment to intravenous amphotericin B (400 mg every 24 h adjusted to the patient's weight). After 7 days, however, the patient developed severe dyspnea requiring oxygen support, aphasia, and general deterioration of health. A second brain scan showed multiple lesions similar to the lesions on the first scan but involving the entire brain parenchyma. The patient died as a result 2 days later. The family did not agree to an autopsy and we were therefore unable to collect brain tissue for microbiologic analysis.

Aspergillus species are members of the eumycetes and are widely distributed in the environment. They are opportunistic pathogens that pose a particular threat to immunosuppressed individuals,^{2,3} particularly those with neutropenia.⁴ The most common species are *Aspergillus fumigatus* and *A. flavus*.³ *Aspergillus* species are ubiquitous in soil and vegetation.² Although aspergillosis mainly affects the lungs,⁵ it can also affect the liver, brain, and skin.³ Between 5% and 27% of invasive aspergillosis cases involve the skin.^{3,6} Cutaneous aspergillosis can be primary or secondary,^{2,7} and these forms can be distinguished by the location and extension of lesions, which

are widespread in secondary infections.^{2,3,7} Secondary cutaneous aspergillosis generally originates from the lungs,¹ but it can also originate from the paranasal sinuses or the upper respiratory tract, although these forms are much less common. Primary aspergillosis is generally caused by direct skin inoculation through wounds from contaminated objects, intravenous lines¹ at venipuncture sites on the arms, or even through dressings covering areas of macerated skin or catheters in patients requiring invasive procedures.⁸

Clinically, aspergillosis manifests as erythematous papules and macules that progress to nodules⁵ and eventually to ulcers with areas of central necrosis.^{3,7} Blisters are uncommon and may, as in our case, contain pus.⁴ Standard diagnostic procedures are the potassium hydroxide technique (or similar) and an incisional skin biopsy with sufficient depth.^{2,7} Histology shows septate hyphae measuring 3 to 5 μm in diameter and 50 to 100 μm in length, 45° branching, absence of blistering with PAS or Gomori methenamine silver stains, and abundant polymorphonuclear cells involving the entire wall, with angiocentric necrosis in many cases.⁴ Serum galactomannan levels should always be tested when aspergillosis is suspected.⁹ A progressive increase to a level over 0.5 in serial measurements points to a diagnosis of invasive bronchopulmonary or systemic aspergillosis, particularly in immunosuppressed patients.⁹ *Aspergillus* infection is confirmed by polymerase chain reaction. Treatment consists of amphotericin B (5 mg/kg/24 h), combined with echinocandins (50–100 mg/d) or voriconazole (200 mg/12 h).¹⁰ Debridement of necrotic lesions and rapid restoration of immunity are important.^{2,7}

We have presented a case of pustular cutaneous aspergillosis in an immunosuppressed patient.




Figure 1 Clinical presentation. A, Painless erythematous nonfluctuant nodule on the left abdominal flank. B, Pustule containing blood-stained pus with a fluid level on an indurated erythematous base.

Download English Version:

<https://daneshyari.com/en/article/8710075>

Download Persian Version:

<https://daneshyari.com/article/8710075>

[Daneshyari.com](https://daneshyari.com)