Accepted Manuscript

Title: Suspected severe acute food protein-induced enterocolitis syndrome caused by cow's milk through breast milk


Author: Ines Vergara Perez, Leticia Vila Sexto

PII: S1081-1206(18)30330-2

DOI: https://doi.org/10.1016/j.anai.2018.04.023

Reference: ANAI 2540

To appear in: Annals of Allergy, Asthma & Immunology



Please cite this article as: Ines Vergara Perez, Leticia Vila Sexto, Suspected severe acute food protein-induced enterocolitis syndrome caused by cow's milk through breast milk, *Annals of Allergy, Asthma & Immunology* (2018), https://doi.org/10.1016/j.anai.2018.04.023.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1	little: Suspected Severe Acute Food Protein-Induced Enterocolitis Syndrome
2	caused by Cow's Milk Through Breast Milk
3	
4	Authors: Ines Vergara Perez MD, Leticia Vila Sexto MD, PhD.
5	Pediatric Allergy Department. Complejo Hospitalario Universitario A Coruña. La
6	Coruña. Spain.
7	
8	Address to correspondence:
9	Leticia Vila Sexto
10	Pediatric Allergy Department. Complejo Hospitalario Universitario A Coruña. La
11	Coruña. Spain.
12	Xubias s/n 15006. A Coruña, Spain
13	e-mail: leticia.vila.sexto@sergas.es; ines.vergara.perez@sergas.es
L 4	Phone number: (+34) 981178688
15	
16	Word count: 933
17	
18	Conflict of interest: none.
19	
20	There was no financial support provided.
21	
22	Key words: food protein-induced enterocolitis, FPIES, cow's milk, food allergy,
23	breast milk.
24	
25	
26	
27	
28	Food protein-induced enterocolitis syndrome (FPIES) is a non-IgE mediated
29	gastrointestinal food hypersensitivity, characterized by profuse vomiting,
30	frequently associated to pallor or/and lethargy that appears within 1 to 3 hours
31	after ingestion of the offending food. Severe cases with protracted emesis
32	develop dehydration, hypotension and acidosis.

Download English Version:

https://daneshyari.com/en/article/8711596

Download Persian Version:

https://daneshyari.com/article/8711596

<u>Daneshyari.com</u>