


Accepted Manuscript

Comparison of allergy immunotherapy medication persistence with a SLIT-tablet vs. SCIT in Germany

Jean-Pierre Allam, MD, Jakob N. Andreasen, MSc, Janina Mette, MSc, Niels Serup-Hansen, MSc, E-MBA, Eike Wüstenberg, MD



PII: S0091-6749(18)30142-8

DOI: [10.1016/j.jaci.2017.12.999](https://doi.org/10.1016/j.jaci.2017.12.999)

Reference: YMAI 13267

To appear in: *Journal of Allergy and Clinical Immunology*

Received Date: 11 January 2017

Revised Date: 27 November 2017

Accepted Date: 4 December 2017

Please cite this article as: Allam J-P, Andreasen JN, Mette J, Serup-Hansen N, Wüstenberg E, Comparison of allergy immunotherapy medication persistence with a SLIT-tablet vs. SCIT in Germany, *Journal of Allergy and Clinical Immunology* (2018), doi: 10.1016/j.jaci.2017.12.999.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comparison of allergy immunotherapy medication persistence with a SLIT-tablet vs. SCIT in Germany

Jean-Pierre Allam MD¹, Jakob N Andreassen MSc², Janina Mette MSc³, Niels Serup-Hansen MSc, E-MBA², Eike Wüstenberg MD^{3,4}

¹Department of Dermatology and Allergy, University of Bonn, Bonn, Germany

²ALK, Hørsholm, Denmark.

³ALK-Abelló, Hamburg, Germany

⁴Department of Otorhinolaryngology, University of Dresden, Dresden, Germany

Corresponding author

Prof. J-P Allam

Department of Dermatology and Allergy

University of Bonn

Bonn, Germany.

Email: Jean-Pierre.Allam@ukb.uni-bonn.de.

Declaration of funding sources: This work was supported by ALK.

Capsule summary

Persistence and adherence for a SLIT-tablet and SCIT were similar in a German population. SLIT discontinuation occurred earlier; patients on SLIT at year 2 were more likely to continue treatment for a third year.

Keywords

SLIT-tablet, SCIT, adherence, medication persistence, grass pollen immunotherapy, retrospective cohort study

Download English Version:

<https://daneshyari.com/en/article/8713205>

Download Persian Version:

<https://daneshyari.com/article/8713205>

[Daneshyari.com](https://daneshyari.com)