


## Accepted Manuscript

Title: Functional outcome of arthroscopic double row repair for Bankart lesion

Authors: Roshan Wade, Bhanuprakash Reddy P V

PII: S0972-978X(18)30115-6  
DOI: <https://doi.org/10.1016/j.jor.2018.03.021>  
Reference: JOR 513



To appear in:

Received date: 12-3-2018  
Accepted date: 19-3-2018

Please cite this article as: Wade R, Reddy P V B, Functional outcome of arthroscopic double row repair for Bankart lesion, *Journal of Orthopaedics* (2018), <https://doi.org/10.1016/j.jor.2018.03.021>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## FUNCTIONAL OUTCOME OF ARTHROSCOPIC DOUBLE ROW REPAIR FOR BANKART LESION

Roshan Wade<sup>1</sup>, Bhanuprakash Reddy P V<sup>2\*</sup>

<sup>1</sup>Assistant professor, Department of Orthopedics, Seth G.S. Medical College and KEM Hospital, Parel, Mumbai, India.

Address: - K/905, Greenwoods, Chakala, Andheri (east), Mumbai-400093, India.

E-mail: - [roshanwade@gmail.com](mailto:roshanwade@gmail.com)

Ph no.: - 98200 50701  
7021 105

<sup>2</sup>Corresponding author

Resident, Department of Orthopedics, Seth G.S. Medical College and KEM Hospital, Parel, Mumbai, India.

Address: - #872/405/49, Brundavana nilaya, 4<sup>th</sup> division, Aishwarya layout, Shanthinagar south, Bychapur road, Devanahalli, Bangalore Rural-562110, India

E-mail: - [dr.bhnauprakash28@gmail.com](mailto:dr.bhnauprakash28@gmail.com)

Ph no.: - 9901574547  
9611734520

### **Abstract**

**Introduction:** - The shoulder joint is the most common major joint to dislocate.

Population aged younger than 20 years, recurrent dislocation rates have been reported to be as high as 90%. For those individuals that continue to experience dislocations surgery is often a good decision. The goal of the Bankart surgery is to reconnect the torn labrum to the glenoid fossa. Double-row Bankart repair can both achieve anatomic reduction and enhance fixation stability. The purpose of our study is to assess the clinical outcome of arthroscopic double row repair.

Download English Version:

<https://daneshyari.com/en/article/8720099>

Download Persian Version:

<https://daneshyari.com/article/8720099>

[Daneshyari.com](https://daneshyari.com)