



Original Article

Length of stay and inpatient mortality trends in primary and revision total joint arthroplasty in the United States, 2000–2014

Matthew Sloan^{a,*}, Neil P. Sheth^b^a Department of Orthopaedic Surgery, University of Pennsylvania, 3737 Market Street, 6th Floor, Philadelphia, PA, 19104, United States^b Department of Orthopaedic Surgery, Pennsylvania Hospital, 800 Spruce Street, 8th Floor, Preston Building, Philadelphia, PA, 19107, United States

ARTICLE INFO

Keywords:

Total joint arthroplasty
Length of stay
Index hospitalization mortality
Comorbidity
Epidemiology

ABSTRACT

Retrospective review of the National Inpatient Sample was performed. LOS and IHM were assessed among TJA patients and assessed by comorbidity status.

LOS among TJA patients decreased from 4.31 to 5.87 days to 2.83–4.49 days. Relative risk for prolonged LOS among high comorbidity patients ranged from 3.01–5.62. IHM per 1,000 was 5.1 for revision THA, 1.8 for revision TKA, 1.1 for primary THA, and 0.4 for primary TKA. Relative risk for IHM in high comorbidity patients ranged from 443.9–780.9 ($p < 0.0001$). LOS and IHM decreased significantly across all TJA procedures. LOS and IHM among the highest comorbidity groups are unacceptably high.

1. Introduction

Total joint arthroplasty (TJA) procedures have increased in volume continuously over the past several decades.^{1–6} Previous studies have projected future volume of primary and revision TJA procedures.^{7,8} However, there has been little discussion of the marked recent changes in length of stay (LOS), index hospitalization mortality and comorbidity prevalence among the primary and revision TJA population since 2000.

Over the past 20 years, the literature has reported on clinical interventions to reduce LOS following TJA, which include improved perioperative pain management using multi-modal therapies,^{9–12} and dedicated clinical pathways focused specifically on the post-operative arthroplasty patient.^{13–16} Recent changes in TJA reimbursement strategies to contain costs, such as bundled payments, demonstrate that pre-discharge costs comprise the majority of expense of care within 30 days of the index procedure.¹⁷ Furthermore, previous studies have demonstrated that reimbursements that financially incentivize shorter LOS will lead to rapid decreases in LOS.¹⁸ Therefore, we expect these recent changes in reimbursement to lead to a decrease in LOS from 2000 to 2014.

Additionally, this study aims to quantify recent changes to safety related to TJA procedures. One study looked at index hospitalization mortality following primary TJA, which has demonstrated declines in recent decades.¹⁹ An earlier study assessed improvement in index hospitalization mortality following implementation of new anesthesia care techniques.²⁰ Cardiac and pulmonary disease history has been associated with 30-day mortality following primary TJA.²¹ We expect

to see similar trends with improved index hospitalization mortality following both primary and revision TJA procedures from 2000 to 2014, in light of modern improvements in anesthesia care and pre-operative screening methods.

Finally, this study aims to address overall changes in the health characteristics of the TJA population. Previous studies have noted increasing prevalence of comorbidity among TJA patients.²² Others have identified that comorbidities, such as obesity and diabetes, portend increased risk of post-operative infection rates and other perioperative complications.²³ We expect to identify increasing prevalence of comorbidity among the TJA population from 2000 to 2014, in line with recent trends.

2. Methods

A retrospective review using discharge data from the National Inpatient Sample and the Nationwide Inpatient Sample (NIS), Healthcare Cost and Utilization Project (HCUP), Agency for Healthcare Research and Quality from 2000 to 2014 was performed.²⁴ This study was given exempt status by the local Institutional Review Board. This database contains 116 million patient records and is a representative sample of discharges among hospitals in the United States. Procedures were identified by International Classification of Diseases, Ninth Revision (ICD-9) codes corresponding to primary total hip arthroplasty (THA; 81.51), primary total knee arthroplasty (TKA; 81.54), revision THA (81.53, 00.70, 00.71, 00.72, 00.73), and revision TKA (81.55, 00.80, 00.81, 00.82, 00.83, 00.84). Procedure volume during the study

* Corresponding author at: Department of Orthopaedic Surgery, University of Pennsylvania, 3737 Market Street, 6th Floor, Philadelphia, PA, United States.
E-mail address: matthew.sloan@uphs.upenn.edu (M. Sloan).

period was evaluated using the appropriate weighting characteristics. Patient population characteristics were evaluated according to discharge variables included in the NIS database corresponding to LOS, index hospitalization mortality, and a comorbidity index score. Comorbidity index was divided into four categories as defined by the 3M All Patient Refined Diagnosis Related Group (APR-DRG) Risk of Mortality subgroups.²⁵ These groups are classified as minor, moderate, major, and extreme likelihood of dying and based on a proprietary algorithm using ICD-9 diagnosis codes. Data for APR-DRGs is available for most patients starting in 2002, and for all patients starting in 2006.

Mean continuous variables were compared between 2000 and 2014 using *t*-test, and proportions were compared by Chi-square test. Linear regression was performed to assess trends in LOS, mortality, and comorbidity prevalence over time. Z-score and relative risk for LOS and mortality by comorbidity subgroup were performed with comparison to “Minor Comorbidity” set as baseline risk. Stata Statistical Software (College Station, TX) was used for all statistical analyses.²⁶

3. Results

3.1. Procedural volume

From 2000 to 2014, TJA procedure volume increased among all categories. Primary THA increased from 160,282 to 371,605 procedures per year. Primary TKA increased from 274,467 to 680,886 procedures per year. Revision THA increased from 34,493 to 50,425 procedures per year. Revision TKA increased from 24,763 to 63,205 procedures per year. During this same time period, TJA population patient characteristics changed markedly. LOS decreased significantly from 2000 to 2014 across all TJA procedures (Fig. 1).

3.2. Length of stay

Primary THA procedure LOS decreased from 4.59 d to 2.83 d (38% decrease, $p < 0.01$), primary TKA LOS decreased from 4.31 d to 2.84 d (34% decrease, $p < 0.01$), revision THA LOS decreased from 5.87 d to 4.49 d (24% decrease, $p < 0.01$), and revision TKA LOS decreased from 4.79 d to 3.80 d (21% decrease, $p < 0.01$). Regression models for all TJA procedure demonstrated a trend of 0.12 d to 0.80 d per year decreases in LOS ($p < 0.01$, R-square > 0.90).

3.3. Index hospitalization mortality

Index hospitalization mortality among all TJA procedures demonstrated a statistically significant 60–70% decrease from 2000 to 2014. Primary THA index hospitalization mortality per 1,000 decreased from 3.5 in 2000 to 1.1 in 2014 (–69%, $p < 0.01$). Primary TKA index

hospitalization mortality decreased from 1.7 in 2000 to 0.43 in 2014 (–75%, $p < 0.01$). Revision THA index hospitalization mortality decreased from 8.8 in 2000 to 5.1 in 2014 (–42%, $p < 0.01$). Revision TKA index hospitalization mortality decreased from 2.3 in 2000 to 1.8 in 2014 (–22%, $p < 0.01$).

3.4. Comorbidity subgroup

Primary and revision TJA procedures differ significantly by comorbidity index ($p < 0.01$). As of 2014, revision THA patients demonstrate the highest proportion of major comorbidity (52%), followed by revision total knee (13%), primary THA (6%), and primary TKA (5%). The THA population demonstrated higher levels of comorbidity compared with the TKA population. Changes in comorbidity subgroup proportions are demonstrated in Fig. 2.

3.5. Comorbidity-stratified length of stay and index hospitalization mortality

LOS and index hospitalization mortality were stratified by comorbidity subgroup. (Fig. 3) LOS increased across comorbidity subgroups in all years. By 2014, LOS ranged 2.55 d to 3.47 d among TJA patients in the minor comorbidity subgroup. Among patients in the extreme comorbidity subgroup, LOS ranged 9.11 d–13.99 d. Revision procedures demonstrated longer LOS compared with primary procedures, and THA procedures demonstrated longer LOS compared with TKA procedures.

Relative risk of prolonged LOS (≥ 5 days) for extreme comorbidity in comparison to minor comorbidity subgroups ranged from 3.01 to 5.62 (95% CI 2.96–5.70, $p < 0.0001$) for primary and revision TJA procedures. (Table 1) The relative risk of extremely prolonged LOS (≥ 10 days) for the extreme comorbidity subgroup ranged from 6.36 to 8.61 (95% CI 6.12–9.02, $p < 0.0001$).

Mortality per 1,000 procedures increased across comorbidity subgroups in all years ($p < 0.01$). (Fig. 3) By 2014, index hospitalization mortality per 1,000 procedures was near 0 for all primary and revision TJA procedures among patients with minor comorbidity. With increasing comorbidity subgroup, index hospitalization mortality per 1,000 procedures increased. Revision THA demonstrated the highest mortality rate in all subgroups. THA procedures demonstrated higher mortality than TKA procedures, and revision procedures demonstrated higher mortality than primary procedures. Index hospitalization mortality for revision THA among patients in the extreme comorbidity subgroup was 160.9 per 1,000 procedures in 2014.

Relative risk of index hospitalization mortality for extreme comorbidity compared with minor comorbidity subgroup ranged from 443.85 to 780.85 (95% CI 301.26–961.63, $p < 0.0001$) among TJA




Fig. 1. Length of stay (LOS) following primary and revision TJA procedures in the United States, 2000–2014.

Download English Version:

<https://daneshyari.com/en/article/8720309>

Download Persian Version:

<https://daneshyari.com/article/8720309>

[Daneshyari.com](https://daneshyari.com)