Accepted Manuscript

Both perforin and FasL are required for optimal CD8 T cell control of autoreactive B cells and autoantibody production in parent-into-F1 lupus mice

Kateryna Soloviova, Maksym Puliaiev, Roman Puliaev, Irina Puliaeva, Charles S. Via

PII: S1521-6616(18)30297-3

DOI: doi:10.1016/j.clim.2018.06.007

Reference: YCLIM 8058

To appear in: Clinical Immunology

Received date: 30 April 2018 Revised date: 18 June 2018 Accepted date: 19 June 2018

Please cite this article as: Kateryna Soloviova, Maksym Puliaiev, Roman Puliaev, Irina Puliaeva, Charles S. Via, Both perforin and FasL are required for optimal CD8 T cell control of autoreactive B cells and autoantibody production in parent-into-F1 lupus mice. Yclim (2018), doi:10.1016/j.clim.2018.06.007

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Both perforin and FasL are required for optimal CD8 T cell control of autoreactive B cells and autoantibody production in parent-into-F1 lupus mice¹

Kateryna Soloviova*, Maksym Puliaiev*, Roman Puliaev*, Irina Puliaeva*, Charles S. Via*

*Pathology Department, Uniformed Services University of Health Sciences, Bethesda, MD 20815

Address correspondence to: Charles S. Via M.D., Department of Pathology,
Room 3B100, 4301 Jones Bridge Road, Uniformed Services University of Health
Sciences, Bethesda, MD 20814. Phone: 301 295 3801; Fax 301 295 1640

Download English Version:

https://daneshyari.com/en/article/8721226

Download Persian Version:

https://daneshyari.com/article/8721226

<u>Daneshyari.com</u>