


Accepted Manuscript

Increased proportion of a CD38^{high}IgD⁺ B cell subset in peripheral blood is associated with clinical and immunological features in patients with primary Sjögren's syndrome

Eriko Ishioka-Takei, Keiko Yoshimoto, Katsuya Suzuki, Ayumi Nishikawa, Hidekata Yasuoka, Kunihiro Yamaoka, Tsutomu Takeuchi



PII: S1521-6616(17)30641-1
DOI: doi:[10.1016/j.clim.2017.10.008](https://doi.org/10.1016/j.clim.2017.10.008)
Reference: YCLIM 7955
To appear in: *Clinical Immunology*
Received date: 1 September 2017
Revised date: 13 October 2017
Accepted date: 19 October 2017

Please cite this article as: Eriko Ishioka-Takei, Keiko Yoshimoto, Katsuya Suzuki, Ayumi Nishikawa, Hidekata Yasuoka, Kunihiro Yamaoka, Tsutomu Takeuchi , Increased proportion of a CD38^{high}IgD⁺ B cell subset in peripheral blood is associated with clinical and immunological features in patients with primary Sjögren's syndrome. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Yclim*(2017), doi:[10.1016/j.clim.2017.10.008](https://doi.org/10.1016/j.clim.2017.10.008)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Increased proportion of a CD38^{high}IgD⁺ B cell subset in peripheral blood is associated with clinical and immunological features in patients with primary Sjögren's syndrome.

Eriko Ishioka-Takei, MD^a, Keiko Yoshimoto, PhD^{a,b,*}, Katsuya Suzuki, MD, PhD^a,
Ayumi Nishikawa, MD, PhD^a, Hidekata Yasuoka, MD, PhD^a, Kunihiro Yamaoka, MD,
PhD^a and Tsutomu Takeuchi, MD, PhD^a

^a *Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine, Tokyo, Japan*

^b *Clinical and Translational Research Center, Keio University Hospital, Tokyo, Japan*

*Correspondence author,

Dr. K. Yoshimoto, Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine, Clinical and Translational Research Center, Keio University Hospital, 35 Shinanomachi, Shinjuku, Tokyo 160-8552, Japan.

E-mail: keikoy@a8.keio.jp Tel/FAX: +81-3-5363-3636

Download English Version:

<https://daneshyari.com/en/article/8721408>

Download Persian Version:

<https://daneshyari.com/article/8721408>

[Daneshyari.com](https://daneshyari.com)