Accepted Manuscript

Response to Pioglitazone in Patients With Nonalcoholic Steatohepatitis With vs Without Type 2 Diabetes

Fernando Bril, MD, Srilaxmi Kalavalapalli, MS, Virginia C. Clark, MD, Romina Lomonaco, MD, Consuelo Soldevila-Pico, MD, I-Chia Liu, Beverly Orsak, RN, Fermin Tio, MD, Kenneth Cusi, MD

 PII:
 S1542-3565(17)31424-6


 DOI:
 10.1016/j.cgh.2017.12.001

 Reference:
 YJCGH 55585

To appear in: *Clinical Gastroenterology and Hepatology* Accepted Date: 1 December 2017

Please cite this article as: Bril F, Kalavalapalli S, Clark VC, Lomonaco R, Soldevila-Pico C, Liu I-C, Orsak B, Tio F, Cusi K, Response to Pioglitazone in Patients With Nonalcoholic Steatohepatitis With vs Without Type 2 Diabetes, *Clinical Gastroenterology and Hepatology* (2018), doi: 10.1016/j.cgh.2017.12.001.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Response to Pioglitazone in Patients With Nonalcoholic Steatohepatitis With vs Without Type 2 Diabetes

Short title: Pioglitazone in NASH and T2DM.

Fernando Bril, MD^{1,2}, Srilaxmi Kalavalapalli, MS¹, Virginia C. Clark, MD³, Romina Lomonaco, MD^{1,2}, Consuelo Soldevila-Pico, MD³, I-Chia Liu¹, Beverly Orsak, RN⁴, Fermin Tio, MD^{5,6}, Kenneth Cusi, MD^{1,2}

¹Division of Endocrinology, Diabetes and Metabolism, University of Florida, Gainesville, Florida. ²Division of Endocrinology, Diabetes and Metabolism, Malcom Randall VAMC, Gainesville, Florida. ³Division of Gastroenterology, Hepatology and Nutrition, University of Florida, Gainesville, Florida. ⁴Diabetes and ⁵Pathology Divisions, University of Texas Health Science Center at San Antonio (UTHSCSA). ⁶Audie L. Murphy Veterans Administration Medical Center (VAMC) San Antonio, Texas.

Financial Support

Supported by the Burroughs Wellcome Fund (K. C.), the American Diabetes Association (1-08-CR-08 [K. C.]), and a VA Merit Award (1 I01 CX000167-01 [K. C.]).

Abbreviations

NASH: nonalcoholic steatohepatitis; T2DM: type 2 diabetes mellitus; NAFLD: nonalcoholic fatty liver disease; IHTG: intrahepatic triglyceride; ALT: alanine aminotransferase; HCC: hepatocellular carcinoma; 1H-MRS: proton magnetic resonance spectroscopy; OGTT: oral glucose tolerance test; DXA: dual-energy absorptiometry; TBF: total body fat; EGP: endogenous glucose production; FFA: free fatty acid; Rd: insulin stimulated glucose disposal; NAS: NAFLD Download English Version:

https://daneshyari.com/en/article/8725085

Download Persian Version:

https://daneshyari.com/article/8725085

Daneshyari.com