Accepted Manuscript

No Significant Association Between Proton Pump Inhibitor Use and Risk of Stroke After Adjustment for Lifestyle Factors and Indication

Long H. Nguyen, Paul Lochhead, Amit D. Joshi, Yin Cao, Wenjie Ma, Hamed Khalili, Eric B. Rimm, Kathryn M. Rexrode, Andrew T. Chan

 PII:
 S0016-5085(17)36710-0

 DOI:
 10.1053/j.gastro.2017.12.006

 Reference:
 YGAST 61581

To appear in: *Gastroenterology* Accepted Date: 11 December 2017

Please cite this article as: Nguyen LH, Lochhead P, Joshi AD, Cao Y, Ma W, Khalili H, Rimm EB, Rexrode KM, Chan AT, No Significant Association Between Proton Pump Inhibitor Use and Risk of Stroke After Adjustment for Lifestyle Factors and Indication, *Gastroenterology* (2018), doi: 10.1053/j.gastro.2017.12.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



No Significant Association Between Proton Pump Inhibitor Use and Risk of Stroke After Adjustment for Lifestyle Factors and Indication

Short Title: Proton Pump Inhibitor Use and Stroke

Authors: Long H. Nguyen^{1,2}, Paul Lochhead^{1,2}, Amit D. Joshi^{1,2}, Yin Cao^{1,2,3}, Wenjie Ma^{1,2}, Hamed Khalili^{1,2}, Eric B. Rimm^{3,4,5}, Kathryn M. Rexrode⁶, Andrew T. Chan^{1,2,5,7}

- 1. Gastrointestinal Unit, Massachusetts General Hospital, Boston, Massachusetts, USA
- 2. Clinical and Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA
- 3. Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA
- 4. Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA
- 5. Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA
- 6. Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA
- 7. Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, Massachusetts, USA

Grant Support: NIH UM1 CA186107 (NHS), NIH UM1 CA167552 (HPFS), NIH R01 HL35464 (EBR), NHLBI R01 HL088521(KMR), and NIH K24 DK098311 (ATC).

Abbreviations: BMI (body mass index), CVD (cardiovascular disease), GERD (gastroesophageal reflux disease), GI (gastrointestinal), H2RA (histamine-2 receptor antagonist), HPFS (Health Professionals Follow Up Study), NHS (Nurses' Health Study), PPI (proton pump inhibitor), PUD (peptic ulcer disease)

Correspondence:

Andrew T. Chan, M.D., M.P.H. Professor of Medicine, Clinical and Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School 55 Fruit Street, GRJ-825C Boston MA 02114, USA <u>achan@partners.org</u>

Disclosures: none Writing Assistance: none

Author Contributions: study concept and design (LHN, ATC); acquisition of data (LHN); analysis and interpretation of data (LHN, PL, ADJ, YC, WM,HK, EBR, KMR, ATC); drafting of the manuscript (LHN); critical revision of the manuscript (EBR, KMR, ATC); statistical analysis (LHN PL, ADJ); study supervision (ATC)

Download English Version:

https://daneshyari.com/en/article/8726736

Download Persian Version:

https://daneshyari.com/article/8726736

Daneshyari.com