


Accepted Manuscript

Title: Critical review of axillary recurrence in early breast cancer

Authors: Carlos A. Castaneda, Pamela Rebaza, Miluska Castillo, Henry H. Gomez, Miguel de La Cruz, Gabriela Calderon, Jorge Dunstan, Jose Manuel Cotrina, Julio Abugattas, Tatiana Vidaurre



PII: S1040-8428(16)30304-3
DOI: <https://doi.org/10.1016/j.critrevonc.2018.06.013>
Reference: ONCH 2579

To appear in: *Critical Reviews in Oncology/Hematology*

Received date: 7-11-2016
Revised date: 20-5-2018
Accepted date: 18-6-2018

Please cite this article as: Castaneda CA, Rebaza P, Castillo M, Gomez HH, de La Cruz M, Calderon G, Dunstan J, Cotrina JM, Abugattas J, Vidaurre T, Critical review of axillary recurrence in early breast cancer, *Critical Reviews in Oncology / Hematology* (2018), <https://doi.org/10.1016/j.critrevonc.2018.06.013>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Critical review of axillary recurrence in early breast cancer

Author names and affiliations

Carlos A. Castaneda^{a,b,*}, Pamela Rebaza^b, Miluska Castillo^b, Henry H. Gomez^a, Miguel de La Cruz^c, Gabriela Calderon^c, Jorge Dunstan^c, Jose Manuel Cotrina^c, Julio Abugattas^c, Tatiana Vidaurre^a.

^aMedical Oncology Department, Instituto Nacional de Enfermedades Neoplasicas, Lima, Peru

^bResearch Department, Instituto Nacional de Enfermedades Neoplasicas, Lima, Peru

^cBreast Cancer Surgery Department, Instituto Nacional de Enfermedades Neoplasicas, Lima, Peru

Contents

1. Introduction
2. Influence of Extension of Axilla Surgery over Recurrence:
 - 2.1. Axillary recurrence rate after Axillary Lymph Node Dissection in breast cancer.
 - 2.2. Axillary recurrence rate after sentinel node biopsy alone in breast cancer with negative sentinel node.
 - 2.3. Axillary recurrence rate after sentinel node biopsy alone in breast cancer with one or two positive sentinel node.
3. Factors related to Axillary Recurrence:
 - 3.1. Predictors of axillary recurrence in breast cancer.
 - 3.2. Prognostic factors after Axillary Recurrence in breast cancer.
4. Management of Axillary Recurrence.
5. Conclusions
 - Conflict of interest
 - Authorship contribution
 - References

***Corresponding author: Carlos A. Castaneda**, Medical Oncology Department, Instituto Nacional de Enfermedades Neoplasicas, Lima, Peru. e-mail: carloscastanedaaltamirano@yahoo.com. Telephone: +51 1 992157220. Fax: + 511 620-4991

Highlights

- Axillary Recurrence (AR) rate after an appropriated treatment in early breast cancer is less than 1%.
- Predictive factors for AR are patient age, HR status, tumor size and histologic grade.
- SLNB is not associated with an increase of AR in SNL-negative and SLN-positive (1-2 nodes) in cT1-2, no palpable nodes cases.
- Prognosis after AR depends on the ER status and the initial systemic treatment.
- Treatment of AR should include systemic therapy for a limited-period.

Download English Version:

<https://daneshyari.com/en/article/8733541>

Download Persian Version:

<https://daneshyari.com/article/8733541>

[Daneshyari.com](https://daneshyari.com)