

Accepted Manuscript

Title: AIDS-associated extensive Kaposi's sarcoma.

Authors: Isaí Medina-Piñón, Martín Wah-Suárez, Luis Arturo Acosta-Calderón, Javier Ramos-Jiménez

PII: S1201-9712(17)30324-7
DOI: <https://doi.org/10.1016/j.ijid.2017.12.012>
Reference: IJID 3120

To appear in: *International Journal of Infectious Diseases*

Received date: 6-11-2017
Revised date: 6-12-2017
Accepted date: 9-12-2017

Please cite this article as: Medina-Piñón Isaí, Wah-Suárez Martín, Acosta-Calderón Luis Arturo, Ramos-Jiménez Javier. AIDS-associated extensive Kaposi's sarcoma. *International Journal of Infectious Diseases* <https://doi.org/10.1016/j.ijid.2017.12.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title: AIDS-associated extensive Kaposi's sarcoma.

Authors: Isaí Medina-Piñón^a, Martín Wah-Suárez^b, Luis Arturo Acosta-Calderón^c, Javier Ramos-Jiménez^a.

Affiliations: a: Infectious Diseases Service. b: Internal Medicine Department. c: Anatomic Pathology and Cytopathology Department. Hospital Universitario "Dr. José E. González" and Medical School. Universidad Autónoma de Nuevo León. Madero y Gonzalitos S/N, Monterrey NL. 64460, México.

Correspondence: Javier Ramos-Jiménez, MD., FIDSA. Head of Infectious Diseases Service, Hospital Universitario "Dr. José E. González". Universidad Autónoma de Nuevo León. México. Madero y Gonzalitos S/N, Monterrey NL. 64460 México. E-mail: javramos31@hotmail.com. Telephone: +52 (81) 83485013.

Highlights

- Kaposi's sarcoma is the most common malignancy in HIV/AIDS patients.
- Histopathologic analysis is the mainstay of workup to confirm diagnosis.
- Gastric Kaposi's sarcoma is the most frequent extracutaneous site.
- Visceral involvement requires combination of HAART and systemic chemotherapy.
- Early diagnosis is essential to avoid advanced presentations.

Keywords: Extensive Kaposi's sarcoma; HIV/AIDS; HHV-8.

Download English Version:

<https://daneshyari.com/en/article/8739096>

Download Persian Version:

<https://daneshyari.com/article/8739096>

[Daneshyari.com](https://daneshyari.com)