

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jmii.com

Brief Communication

Nasal *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* carriage among college student athletes in northern Taiwan

Hong-Kai Wang^{a,c}, Chun-Yen Huang^{a,c}, Chih-Jung Chen^{a,b},
Yhu-Chering Huang^{a,b,*}

^a School of Medicine, Chang Gung University, Gueishan, Taoyuan, Taiwan

^b Department of Pediatrics, Chang Gung Memorial Hospital at Linkou, Gueishan, Taoyuan, Taiwan

Received 13 July 2016; received in revised form 18 October 2016; accepted 7 November 2016

Available online ■ ■ ■

KEYWORDS

Methicillin-resistant
Staphylococcus aureus;
Community-associated;
Athlete;
College student;
Taiwan

Abstract Of 259 college students in northern Taiwan surveyed, nasal carriage rate of *Staphylococcus aureus* and methicillin-resistant *S. aureus* (MRSA) was 22.4% and 1.54%, respectively and no significant difference was found between athlete students and non-athlete students. Three of four MRSA isolates belonged to sequence type 59, the endemic community clone. Copyright © 2017, Taiwan Society of Microbiology. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Staphylococcus aureus, including methicillin-resistant (MRSA), is an important pathogen of serious infections in

humans.¹ Not only a hospital pathogen, MRSA is also a pathogen circulated in the community since 1990s.^{1,2} MRSA can colonize on anywhere of human bodies, particularly in the anterior nares.³ The carriage of *S. aureus*, including MRSA, is a significant risk factor for subsequent infection.^{1–3} Community-associated (CA)-MRSA isolates, initially identified in pediatric populations,^{1,2} were also reported in certain adult population which included competitive sports participants,^{1,2} especially the sports with frequent body contacts. There have been scanty reports on nasal MRSA carriage among competitive sports

* Corresponding author. Department of Pediatrics, Chang Gung Memorial Hospital, No. 5, Fu-Shin Street, Gueishan, 333, Taoyuan, Taiwan. Fax: +886 3 3288957.

E-mail address: ychuang@adm.cgmh.org.tw (Y.-C. Huang).

^c First two authors equally contributed to this work.

<http://dx.doi.org/10.1016/j.jmii.2016.11.005>

1684-1182/Copyright © 2017, Taiwan Society of Microbiology. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Wang H-K, et al., Nasal *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* carriage among college student athletes in northern Taiwan, Journal of Microbiology, Immunology and Infection (2017), <http://dx.doi.org/10.1016/j.jmii.2016.11.005>

participants,^{4–8} and none from Taiwan. Therefore, we conducted this study to elucidate the carriage rate of *S. aureus*, including MRSA, among competitive sports participants in Taiwan.

Methods

This study was approved by Institutional Reviewing Board of Chang Gung Memorial Hospital (reference No: 102-1946B) and a written informed consent was obtained from each participant. Between June and September, 2013, 121 students from Chang Gung University (CGU) (non-athletes group) and 139 students of the department of sport training science from National Taiwan Sports University (NTSU) (athletes group), respectively, were recruited. Athlete students were further categorized into two groups by with or without close body contact. The contact sport group included basketball, Judokas, Taekwondo, and soccer. The non-contact group included tennis and table tennis. Eligible subjects were invited to participate, and specimens were obtained from their anterior nares for the detection of *S. aureus*. A questionnaire was completed for each subject, including demographics and potential risk factors for the acquisition of *S. aureus* and MRSA. Subjects whose roommates, close friends, or team mates were smokers were categorized as frequent smoking exposure.

A nasal swab specimen was collected from the anterior nares for each subject by a dry Copan Transystem Liquid Stuart swab (Venturi Transystem, Copan Innovation Ltd, Limerick, Ireland). Each nasal swab was circled in the participant's both nasal vestibules. The swabs were placed into the transport media and transported to the laboratory within 4 h. *S. aureus* as well as MRSA was identified and the antimicrobial susceptibility to 10 antibiotics was determined by using the disk-diffusion method, according to the recommendations of CLSI.

For molecular characterization, all *S. aureus* isolates were characterized by pulsed-field gel electrophoresis (PFGE), staphylococcal cassette chromosome (SCC*mec*) typing (only for MRSA), and detection of Panton–Valentine leukocidin (PVL) genes. Some isolates of representative pulsotypes were further typed by multilocus sequence type (MLST) and *spa* gene typing. All the molecular methods as well as SCC*mec* typing for type V_T were described elsewhere.⁹

Results

A total of 259 students, including 139 athlete students and 120 non-athlete students, were recruited. Age ranged from 18 to 35 years, with a mean of 20.6 years. In the athletes group, the mean age was 19.7 years and in the non-athletes group, 21.5 years. The male to female ratio was 1:0.43. Of the athletes group, 73% participated in contact sports and 27% non-contact sports.

Overall, *S. aureus* was identified in 58 students (22%), including MSSA in 54 students (21%) and MRSA in 4 students (1.5%). The mean ages of carriers of *S. aureus* and MRSA were 20.8 ± 2.89 years and 19.0 ± 1.16 years, respectively. No significant difference was found between the two groups (athletes vs. non-athletes) in terms of nasal carriage of

MSSA (21% vs. 21%) and MRSA (0.72% vs. 2.50%). Among the athlete students, *S. aureus* carriage rate varied from 10% for the Taekwondo category to 40% for the soccer category (Table 1).

Since only four students had MRSA colonization, it could not be calculated in an effective method to identify the risk factors. We analyzed the risk factor of carrying *S. aureus* and MSSA. The comparison of potential risk factors for the carriage of *S. aureus* between carriers and non-carriers is shown in Supplementary Table. Household members working as health care workers (HCWs) was a risk factor of carrying *S. aureus* ($p = 0.033$). Smoking or frequent smoking contact was a marginal protective factor ($p = 0.065$). The carriage rate for non-contact sports group was higher than that for contact sports group (32% vs. 18%, $p = 0.061$). There was no statistically significant difference between the carriers and non-carriers of *S. aureus* in terms of age, gender, prior hospitalization, using artificial catheters/tubes.

Fig. 1 reveals the detailed molecular characteristics of the four MRSA isolates. Three PFGE patterns were identified. All four isolates carried SCC*mec* IV or V_T. None of the isolates carried PVL genes. Three isolates from non-athletes belonged to sequence type (ST) 59/*spa* t437, the endemic CA-MRSA strains in Taiwan.

Of 54 MSSA isolates, a total of 18 pulsotypes were identified, nine pulsotypes identified in non-athlete students (unidentified pattern for one isolate), and 16 pulsotypes in athlete students (Table 1). Pulsotype BA, accounting for 15 (28%) of the MSSA isolates, was the dominant type and prevailed in both universities. The 15 isolates of pulsotype BA belonged to at least three STs, namely ST 6, 7 and 97, and at least six *spa* types. None of the isolates carried PVL genes. All 54 MSSA isolates were susceptible to vancomycin, teicoplanin, linezolid, trimethoprim-sulfamethoxazole and ciprofloxacin. Five of eight isolates which were resistant to erythromycin were also resistant to clindamycin. 39 isolates (72%) were resistant to penicillin. No statistically significant difference was found between the isolates from athletes and non-athletes in terms of antibiotic resistance rates.

Discussion

Results from the present study indicated that 1.54% of college students in northern Taiwan, 2.50% for non-athletes group and 0.72% for athletes group, harbored MRSA in their nares, and there was no statistically significant difference in carriage rate of MRSA between the two groups. The carriage rate of MRSA among the two universities' students was lower than that for general population (3.8%)^{1,2} and healthcare workers (5.0–7.8%)^{1,2} in northern Taiwan but was consistent with that for medical students (2.2%).²

Reviewing MRSA colonization among competitive sport participants published in the English language literature,^{4–8} we found that the carriage rate ranged from 0% to 37%. In the report from Ohio, USA, none of the recruited competitive sport participants harbored MRSA.^{4,7} Another report from Tennessee, USA,⁵ showed MRSA carriage rate varied significantly through the athletic season, from 4% to 23%, with a peak during the time of highest athletic activity. We

Download English Version:

<https://daneshyari.com/en/article/8741014>

Download Persian Version:

<https://daneshyari.com/article/8741014>

[Daneshyari.com](https://daneshyari.com)