Accepted Manuscript

Quantitative measurement of cartilage volume is possible using two-dimensional magnetic resonance imaging data sets

L.F. Schaefer, V. Nikac, J.A. Lynch, J. Duryea

PII: \$1063-4584(18)31147-6

DOI: 10.1016/j.joca.2018.04.005

Reference: YJOCA 4215

To appear in: Osteoarthritis and Cartilage

Received Date: 15 September 2017

Revised Date: 9 April 2018
Accepted Date: 11 April 2018

Please cite this article as: Schaefer LF, Nikac V, Lynch JA, Duryea J, Quantitative measurement of cartilage volume is possible using two-dimensional magnetic resonance imaging data sets, *Osteoarthritis and Cartilage* (2018), doi: 10.1016/j.joca.2018.04.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Quantitative measurement of cartilage volume is possible using two-dimensional magnetic resonance imaging data sets

L. F. Schaefer¹, V. Nikac¹, J.A. Lynch², J. Duryea¹

¹Brigham and Women's Hospital, Harvard Medical School. Boston, MA

²University of California, San Francisco. San Francisco, CA

Keywords: Osteoarthritis, Cartilage, Knee, Magnetic Resonance Imaging, Segmentation Software

Address for correspondence:

Lena Schaefer MD

Department of Radiology,

Brigham and Women's Hospital, Harvard Medical School,

75 Francis Street, Boston, MA 02115

lenafranziskaschaefer@yahoo.com

Download English Version:

https://daneshyari.com/en/article/8741572

Download Persian Version:

https://daneshyari.com/article/8741572

<u>Daneshyari.com</u>