Accepted Manuscript



Two extracellular sialidases from *Bifidobacterium bifidum* promote the degradation of sialyl-oligosaccharides and support the growth of *Bifidobacterium breve*

Keita Nishiyama, Aki Nagai, Kazuya Uribayashi, Yuji Yamamoto, Takao Mukai, Nobuhiko Okada

PII: S1075-9964(18)30089-1

DOI: 10.1016/j.anaerobe.2018.05.007

Reference: YANAE 1888

To appear in: *Anaerobe*

Received Date: 27 February 2018

Revised Date: 17 May 2018

Accepted Date: 18 May 2018

Please cite this article as: Nishiyama K, Nagai A, Uribayashi K, Yamamoto Y, Mukai T, Okada N, Two extracellular sialidases from *Bifidobacterium bifidum* promote the degradation of sialyl-oligosaccharides and support the growth of *Bifidobacterium breve*, *Anaerobe* (2018), doi: 10.1016/j.anaerobe.2018.05.007.

Anaerobe

Two extracellular sialidases from *Bifidobacterium bifidum* promote the degradation of sialyl-oligosaccharides and support the growth of *Bifidobacterium breve*

Keita Nishiyama[a*], Aki Nagai[a], Kazuya Uribayashi[a], Yuji Yamamoto[b], Takao Mukai[b], Nobuhiko Okada[a]

[a]Department of Microbiology, School of Pharmacy, Kitasato University, Minato-ku, Tokyo, Japan

[b]Department of Animal Science, School of Veterinary Medicine, Kitasato University, Towada, Aomori, Japan

[*]Corresponding author.

Tel.: +81-3-5719-6256; Fax: +81-3-3444-4831

E-mail: nishiyamak@pharm.kitasato-u.ac.jp

Declarations of interest: none.

Abbreviations: HMO, human milk oligosaccharides; PCM, porcine colonic mucin; TLC, thin-layer chromatography; 6′SL, 6′ sialyllactose