



ELSEVIER

Contents lists available at ScienceDirect

## Journal of Infection and Public Health

journal homepage: <http://www.elsevier.com/locate/jiph>



# A multi-faceted approach of a nursing led education in response to MERS-CoV infection<sup>☆</sup>

Jaffar A. Al-Tawfiq<sup>a,b,\*</sup>, Siobhan Rothwell<sup>c</sup>, Heather A. McGregor<sup>c</sup>, Zeina A. Khouri<sup>c</sup>

<sup>a</sup> Speciality Medicine Division, Johns Hopkins Aramco Healthcare, Dhahran, Saudi Arabia

<sup>b</sup> Indiana University School of Medicine, Indianapolis, IN, USA

<sup>c</sup> Nursing Department, Johns Hopkins Aramco Healthcare, Dhahran, Saudi Arabia

### ARTICLE INFO

#### Article history:

Received 23 May 2017

Accepted 4 August 2017

#### Keywords:

Coronavirus  
Education  
MERS-CoV  
Middle East respiratory syndrome  
Saudi Arabia  
Outbreak preparedness  
Hospital care

### ABSTRACT

**Background:** The emergence of the Middle East respiratory syndrome–coronavirus (MERS-CoV) resulted in multiple healthcare associated outbreaks. In response, we developed a nurse-led program to screen and triage patients with MERS-CoV infection.

**Methods:** A formal educational program was implemented to ensure a standardized approach to care planning and delivery. The essential skills that were included were the use of Personal Protective Equipment (PPE) including gown, gloves, head cover and N95 mask, hand hygiene, the practice of donning and doffing and the collection of MERS-CoV nasopharyngeal specimens.

**Results:** A core group of nurses were identified from high-risk units. The education program consisted of four skill-days. Nurses were assigned a pre-course work from Mosby's Nursing Skills, an online skill module, donning and doffing of PPE, and the process for obtaining a nasopharyngeal swab. The skill lab incorporated multiple methods such group discussion, watching an on-line video, and a simulated demonstration and practice on a mannequin. In total, 450 nurses attended the Nasopharyngeal Skills Day and 1000 nurses received training. Donning and doffing PPE and N95 mask fit testing has become an annual mandatory competency requirement for staff in nursing and clinical services.

**Conclusion:** The application of specified protocols minimizes the risks of cross infection, placing emphasis on patient and staff safety as well as expediting the patient to definitive treatment. Nurse educators have an instrumental role in training nurses and other healthcare providers on the specifics needed to identify, contain and manage patient presenting with MERS-CoV.

© 2017 The Authors. Published by Elsevier Limited on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Introduction

The emergence of the Middle East Respiratory Syndrome Coronavirus (MERS-CoV) in the Kingdom of Saudi Arabia (KSA) in 2012 [1] resulted in multiple healthcare associated infections in Al-Hasa, Jordan, Jeddah, Taif, and the Republic of Korea [2–9]. A coordinated response is necessary to minimize the risks to healthcare providers and other patients. It was found that close contact, such as providing unprotected care to a patient can result in person to person transmission. Simple infection control measures result of the first outbreak in Al-Hasa, KSA [2]. MERS-CoV as other emerging infec-

tious diseases had been associated with fears and anxiety of taking care of patients [10,11]. Thus, we developed a nurse-led program to stream line the screening process and to establish a cohort of healthcare workers to obtain nasopharyngeal swabs for MERS-CoV testing. In this report, we describe the program and highlight its salient components.

### Materials and methods

The study was conducted at Johns Hopkins Aramco Healthcare with a multifaceted approach.

### Intervention

The hospital response included a review of international guidelines [12–14] and those offered by the Saudi Arabian Ministry of Health [15]. Advice and support was provided by the Infection Control and Prevention Department with regards to disseminating

<sup>☆</sup> The study was approved by the IRB of the Johns Hopkins Aramco Healthcare, #17-13.

\* Corresponding author at: P.O. Box 76, Room A-428-2, Building 61, Dhahran Health Center, Johns Hopkins Aramco Healthcare, Dhahran 31311, Saudi Arabia. Fax: +966 13 877 3790.

E-mail addresses: [jaffar.tawfiq@jhah.com](mailto:jaffar.tawfiq@jhah.com), [jaltawfi@yahoo.com](mailto:jaltawfi@yahoo.com) (J.A. Al-Tawfiq).

<http://dx.doi.org/10.1016/j.jiph.2017.08.006>

1876-0341/© 2017 The Authors. Published by Elsevier Limited on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Table 1**  
The skills labs incorporated multiple skills.  
(PPE: Personal Protective Equipment)

Skills	Method
PPE Donning and Doffing	Demonstration
Correct use of N 95 mask	Demonstration
Orientation to Laboratory and MOH forms required for nasopharyngeal specimen testing	Discussion
Process for transportation of nasopharyngeal swab to Laboratory for testing	Discussion
Instruction on how to obtain a nasopharyngeal swab	Procedure on-line video obtained from the New England Journal of Medicine [18]
Obtaining a nasopharyngeal swab from a patient	Simulated demonstration and practice on a mannequin – competency assessed



**Fig. 1.** Nasopharyngeal Training Station: each nurse joined a group where a nurse educator simulated the procedure, using a mannequin.

information and developing processes to assist an interdisciplinary response team. It became obvious that good infection control and prevention practices would be necessary to not only care for patients who had tested positive for MERS-CoV but also for suspected cases and to protect staff in the delivery of the required care. Thus, a new process had to be implemented in order to reduce the risk of transmission of MERS-CoV. The study was approved by the Institutional Review Board.

### Education

Staff awareness of MERS-CoV symptoms and initial management was the first step. Screening and subsequent isolation of patients presenting to the Emergency Department or Primary Care with symptoms suggestive of MERS-CoV would reduce the risks of transmission. Alerting staff to the signs and symptoms aimed to increase early recognition of the virus in patients presenting to the hospital. The next stage required a formal educational strategy to ensure a standardized approach to care planning and delivery. Following local and international guidelines [12–14], the Clinical Practice, Education, Research and Education Division (CPERD) commenced the education and training of all nursing staff to ensure competency in caring for patients with actual or suspected MERS-CoV. It was recognized the hospital response would need to include more advanced infection control measures in order to ensure the protection of health care providers. Recommendations from the CDC included employing standard, contact and airborne precautions which was used to devise the education and training program [12–15]. The essential skills that were included in the program included: the use of Personal Protective Equipment (PPE) including gown, gloves, head cover and N95 mask, hand hygiene, the practice of donning and doffing and the collection of MERS-CoV nasopharyngeal specimens.

### Nasopharyngeal Swab Educational Program

Nasopharyngeal swab specimens are used to obtain samples for MERS-CoV testing [16]. Therefore a good technique is essential to

ensure that an adequate sample was obtained. It was suggested at that time that lower respiratory tract samples should be obtained for confirmation of the MERS-CoV virus [17]. Lower respiratory samples were obtained whenever this was possible.

### Results

A core group was identified from the Emergency Department, Primary Care, the dedicated Isolation Unit, and the Hemodialysis Unit. The education program consisted of four skill-days consisting of one hour rolling sessions. The program was then extended to other wards and units who may admit or take care of MERS-CoV patients. Nurses were assigned a pre-course work from Mosby's Nursing Skills, an online skill module, donning and doffing of PPE, and the process for obtaining a nasopharyngeal swab. The skill lab incorporated multiple methods such as demonstration, group discussion, watching an on-line video obtained from the New England Journal of Medicine [18], and a simulated demonstration and practice on a mannequin (Table 1).

Nurses were re-orientated to the correct process of donning and doffing PPE including N95 mask through a demonstration process. The nurses were then shown a video developed by the New England Journal of Medicine entitled: Procedure – Collection of Nasopharyngeal Specimens with the Swab Technique [18]. Each nurse joined a skills table where a nurse educator simulated the procedure, using a mannequin, to obtain a nasopharyngeal swab (Fig. 1). Special consideration was given to the position of the patient, the depth the swab was to be placed and the length of time the swab was to be in contact with the mucosal surface. The procedure was practiced until all nurses were comfortable with the process. The skills lab also addressed specific requirements outlined by the Prevention and Control of Infection Committee, the hospital Laboratory and the Ministry of Health including the necessary forms (Fig. 2), transportation of the specimens to the Laboratory and the notification requirements.

The reporting forms were revised by the educators to ensure all required components were documented. The process for notifying the hospital Epidemiology and the Ministry of Health was updated

Download English Version:

<https://daneshyari.com/en/article/8746819>

Download Persian Version:

<https://daneshyari.com/article/8746819>

[Daneshyari.com](https://daneshyari.com)