



Alexandria University Faculty of Medicine
Alexandria Journal of Medicine

<http://www.elsevier.com/locate/ajme>



CASE REPORT

Isolated cutaneous leishmaniasis over face – A diagnostic dilemma

Santosh K. Swain^{a,*}, Ishwar C. Behera^b, Mahesh C. Sahu^c, Maitreyee Panda^d

^a Department of Otorhinolaryngology, IMS and SUM Hospital, Siksha “O” Anusandhan University, K8, Kalinganagar, Bhubaneswar 751003, Odisha, India

^b Department of Community Medicine, IMS and SUM Hospital, Siksha “O” Anusandhan University, K8, Kalinganagar, Bhubaneswar 751003, Odisha, India

^c Central Research Laboratory, IMS and SUM Hospital, Siksha “O” Anusandhan University, K8, Kalinganagar, Bhubaneswar 751003, Odisha, India

^d Department of Dermatology, IMS and SUM Hospital, Siksha “O” Anusandhan University, K8, Kalinganagar, Bhubaneswar 751003, Odisha, India

Received 29 August 2015; revised 15 December 2015; accepted 16 December 2015

KEYWORDS

Cutaneous leishmaniasis;
 Fungal dermatitis;
 Leprosy;
 Diagnostic dilemma

Abstract Cutaneous Leishmaniasis (CL) is a disease caused by an intracellular protozoa belong to the genus Leishmania, transmitted by the bite of a sandfly. It has diverse clinical presentation and may create a public health problem in endemic countries. CL is often confused with lepromatous leprosy, pimples and fungal dermatitis. This case is an isolated cutaneous variety in facial region which was mistaken and treated initially for fungal dermatitis and then for leprosy by local physicians. Smears examined from the skin lesion confirmed Leishmania amastigotes. The isolated localized CL may create confusion and its many differential diagnoses made delaying in the diagnosis.

© 2015 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Leishmaniasis is a vector borne disease and caused by a kinetoplastid protozoan parasite. The parasite is transmitted from one patient to another through the bites of female sandfly, or occasionally through non-vector routes including congenital, blood transfusion, sexual, laboratory acquired and person-

to-person.¹ Leishmaniasis can be divided into three forms varying in severity from spontaneously healing dermatological ulcers in Cutaneous leishmaniasis (CL) and destructive form of mucocutaneous leishmaniasis to fatal form of visceral leishmaniasis (VL). CL is seen throughout Africa, South America, Asia, the Middle East, and the Mediterranean regions. It has diverse clinical manifestations, which poses a public health problem in endemic countries.² This disease can affect the skin, viscera or mucocutaneous area. Often CL cases are mistaken for common dermatological disorders such as fungal infections, bacterial infections, eczema or chronic ulcers that may delay in diagnosis and treatment. In some cases, patients consult several doctors before CL diagnosis. This situation leads to delay in the exact diagnosis and inadvertent exposure to

* Corresponding author at: Department of Otorhinolaryngology, IMS and SUM Hospital, Bhubaneswar, Odisha, India. Cell: +91 9556524887.

E-mail address: santoshvoltaire@yahoo.co.in (S.K. Swain).

Peer review under responsibility of Alexandria University Faculty of Medicine.

<http://dx.doi.org/10.1016/j.ajme.2015.12.003>

2090-5068 © 2015 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

adverse events of inappropriate treatment. Here an isolated lesion over the face made a misdiagnosis for fungal dermatitis initially and then leprosy by local physicians for which patient had taken prolonged unnecessary treatment due to delay in diagnosis.

2. Case report

A 48 year old male from northeast part of India presented to Outpatient department of Otorhinolaryngology for non-healing skin lesions over the face (Fig. 1A and B). From history, it was found that he was treated by many physicians by multiple drugs since two years, initially for fungal dermatitis and then for leprosy but lesions were not healed. On examination there are several small painless papular lesions with indurations. The papular lesions are showing central ulceration with erythematous borders over the forehead and cheek. He had no significant medical history and patient was in good health. He had no history of fever, lymphadenopathy, hepatosplenomegaly or weight loss which are typical of VL. There were no signs of neuropathy. General clinical examinations and routine blood investigations were within normal limit. HIV test was negative. Slit skin smears from facial lesions were stained with Ziehl-Nielsen stain and revealed negative for acid-fast bacilli.

The biopsies and smears were taken from the facial lesions under the local anaesthesia and sent for histopathological examination. Light microscopy revealed skin with hyperkeratosis, acanthosis and parakeratosis. The dermis was invaded with large pink coloured histiocytes and chronic inflammatory cells. The histiocytes contained inside dot-like organisms typical of cutaneous leishmania (Fig. 2). He was treated with sodium stibogluconate for 3 weeks (intramuscular 20 mg/kg daily for 3 weeks). The cutaneous lesions of the face disappeared and there has been no evidence of recurrence after two years.

3. Discussion

Leishmaniasis is caused by unicellular, flagellate, intracellular protozoa belonging to the genus *Leishmania* transmitted from

animals to humans through phlebotomine sandfly vectors. Common species of *Leishmania* responsible for CL in old world are *L. major* and *L. tropica* whereas *L. mexicana*, *L. panamenesis* and *L. braziliensis* in new world. The geographical distribution of Leishmaniasis is confined to temperate and tropical countries, the living area of the sandfly. Ninety per centages of cases of CL occur in Brazil, Afghanistan, Iran, Algeria, Peru, and Syria whereas visceral leishmaniasis is found in Bangladesh, India, Brazil, Nepal and Sudan.³ In India, indigenous cases of CL are mainly confined to hot dry northwestern region and are endemic in the western Thar desert of Rajasthan.⁴ Few cases have been observed in the states such as Kerala, Assam and Haryana.⁵ Apart from this it was documented in northwestern part of Indo-Gangetic plain, along with dry areas of Indo-Pakistan border from Amritsar to Gujarat.⁶ In southern part of India, CL has been reported from Malappuram and Trivandrum of Kerala.⁷ The CL is usually caused by *L. tropica* and man is the most common reservoir in India.

Leishmaniasis has been considered tropical afflictions and constitutes one of the six entities in the World health organization/Tropical disease research (WHO/TDR) list of the most important diseases.^{8,9} Leishmaniasis is endemic in 88 countries of the five continents with total of 350 million people at risk and heaves 12 million cases. Out of the 88 endemic countries, 22 are in the New World and 66 in the Old World with an estimated incidence of 1–1.5 million cases of CL and 5 lakhs cases of visceral leishmaniasis (VL).⁹ Even it has widespread geographical distribution, human leishmaniasis is often very focal within an endemic region, leading to 'hotspots' of the disease transmission.⁵ Localized cutaneous lesions of leishmaniasis may resemble to other skin conditions such as lepromatous leprosy, cutaneous anthrax, blastomycosis, sporotrichosis, eczema, fungal skin infections, Mycobacterium marinum infections, sarcoidosis and infected insect bites. Isolated cutaneous lesion of leishmaniasis over the face may mimic lupus vulgaris, chronic discoid erythematosus, cutaneous lymphoma and erysipelas.^{10,11} Infections by *Leishmania* are increasing worldwide due to tourism and job related travel and refugees.¹²

The parasite multiplies inside the macrophages and other types of reticuloendothelial cells will rupture and release the



Figure 1 Cutaneous leishmaniasis lesions seen over the face (A) frontal view and (B) lateral view.

Download English Version:

<https://daneshyari.com/en/article/8752347>

Download Persian Version:

<https://daneshyari.com/article/8752347>

[Daneshyari.com](https://daneshyari.com)