Accepted Manuscript

A high anti-cholinergic burden is associated with a history of falls in the previous year in middle-aged women: findings from the Aberdeen Prospective Osteoporosis Screening Study

Andrew D. Ablett, Adrian D. Wood, Rebecca Barr, Jordan Guillot, Alison J. Black, Helen M. Macdonald, David M. Reid, Phyo K. Myint

PII: S1047-2797(17)30933-X

DOI: 10.1016/j.annepidem.2018.05.011

Reference: AEP 8410

To appear in: Annals of Epidemiology

Received Date: 19 September 2017

Revised Date: 14 May 2018 Accepted Date: 26 May 2018

Please cite this article as: Ablett AD, Wood AD, Barr R, Guillot J, Black AJ, Macdonald HM, Reid DM, Myint PK, A high anti-cholinergic burden is associated with a history of falls in the previous year in middle-aged women: findings from the Aberdeen Prospective Osteoporosis Screening Study, *Annals of Epidemiology* (2018), doi: 10.1016/j.annepidem.2018.05.011.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

A high anti-cholinergic burden is associated with a history of falls in the previous year in middle-aged women: findings from the Aberdeen Prospective Osteoporosis Screening Study

Andrew D Ablett^{1,2}, Adrian D Wood^{1,2}, Rebecca Barr^{2,3,5}, Jordan Guillot¹, Alison J Black^{2,4}, Helen M Macdonald^{2,3}, David M Reid^{2,3}, and Phyo K Myint^{1,2,4}

Keywords: Anti-cholinergic, epidemiology, bone mineral density, falls, fracture, women, middle age

Word count: 2855

Table 1: Participant characteristics by anti-cholinergic burden score in 3,883 middle-aged women of APOSS at second visit.

Table 2: Results of logistic regression analysis examining the association between anticholinergic burden (reference category = 0) a history of falls over the previous 12 months.

Table 3: Results of logistic regression analyses examining association between anticholinergic burden (reference category = 0) and hip and spine bone mineral density.

Supplementary Table: Anti-cholinergic burden scoring table.

Correspondence to:

Professor Phyo Kyaw Myint Room 4:013, Polwarth Building, School of Medicine, Medical Sciences and Nutrition, Foresterhill, Aberdeen, AB25 2ZD, Scotland, UK. Tel: +44 (0) 1224 437841, Fax: +44 (0) 1224 43791

Mail to: phyo.myint@abdn.ac.uk

¹Ageing Clinical & Experimental Research (ACER) Team, Institute of Applied Health Sciences, School of Medicine, Medical Sciences, and Nutrition, University of Aberdeen, UK ²Aberdeen Centre for Arthritis and Musculoskeletal Health, University of Aberdeen, UK ³Arthritis and Musculoskeletal Medicine Research Programme, Institute of Medical Sciences, University of Aberdeen, UK

⁴NHS Grampian, Foresterhill Road, Foresterhill, Aberdeen, UK

⁵Medicines Monitoring Unit, University of Dundee, Dundee, UK

Download English Version:

https://daneshyari.com/en/article/8753201

Download Persian Version:

https://daneshyari.com/article/8753201

<u>Daneshyari.com</u>