



Contents lists available at ScienceDirect

European Journal of Internal Medicine

journal homepage: www.elsevier.com/locate/ejim

Anticipating care needs of patients after discharge from hospital: Frail and elderly patients without physiological abnormality on day of admission are more likely to require social services input

Subbe C.P.^{a,b,*}, Goulden N.^c, Mawdsley K.^c, Smith R.^d

^a Acute, Respiratory & Critical Care Medicine, Ysbyty Gwynedd, United Kingdom

^b School of Medical Sciences, Bangor University, Bangor LL57 2AS, United Kingdom

^c North Wales Organisation for Randomised Trials in Health, School of Healthcare Sciences, Bangor University, Bangor LL57 2PZ, United Kingdom

^d Wrexham County Borough Council, Guildhall, Wrexham LL11 1AY, United Kingdom

ARTICLE INFO

Article history:

Received 15 August 2017

Received in revised form 13 September 2017

Accepted 23 September 2017

Available online xxxxx

Keywords:

Frailty

Hospital

Patient discharge

Social work

Triage

ABSTRACT

Introduction: Acute admissions to hospital are rising. As a part of a service evaluation we examined pathways of patients following hospital discharge depending on data available on admission to hospital.

Methods: We merged data available on admission to the Wrexham Maelor hospital from an existing data-base in the Acute Medical Unit with follow up data from local social services as part of a data sharing agreement. Patients requiring support by social services post-discharge were matched with patients not requiring social services from the same post-code.

Results: Stepwise logistic regression analysis identified candidate variables predicting likely support need. Decision tree analysis identified sub-groups of patients with higher likelihood to require support by social services after discharge from hospital. We found patients with normal physiology on admission as evidenced by a value of zero for the National Early Warning Score who were frail or older than 85 years were most likely to require support after discharge.

Conclusions: Information available on admission to hospital might inform long term care needs. Prospective testing is needed. The algorithms are prone to be dependent on availability of local services but our methodology is expected to be transferable to other organizations.

© 2017 Published by Elsevier B.V. on behalf of European Federation of Internal Medicine.

1. Introduction

Frail elderly patients often get admitted with seemingly minor illnesses, yet end up staying in hospital for a long time. Part of the reason for this is that rehabilitation often starts late during the course of the hospital stay, at a time when patients have already lost significant strength and confidence. Early recognition and treatment of frailty is therefore a concept that might reduce length of hospital stay and increase patients' independence and satisfaction.

Acute Medicine is the port of entry to UK hospitals for 50% of patients and an even higher proportion of the frail elderly [1]. Frailty, if not detected and supported in a timely fashion leads to prolonged hospital stay, higher levels of dependency and higher mortality [2,3].

In 2014 we showed with the help of a SHINE grant from the Health Foundation that a navigator supported triage system based

on parameters available on admission to hospital can drive selection of patients for early discharge and reduce hospital length of stay [4]. Data from a small subset of patients suggested that time to referral for intermediate care services was cut by 14 days with the usage of the Clinical Frailty Scale (CFS) [5] without an increase in futile referrals.

Our previous research did not include follow-up after discharge from hospital.

In order to improve the patients experience of hospital and to optimize usage of resources we hypothesise that it is possible to recognize frailty on admission to hospital, that this recognition will lead to earlier referral to the multi-disciplinary intermediate care or rehabilitation team and therefore to earlier discharge to patients' own homes with a greater degree of independence.

Fy Nhaid (Frailty: timely recognition of Need for Home support After In-patient Discharge) aimed to develop locally applicable simple criteria for earlier recognition of patients at risk of increased care needs post-hospital discharge that allow prospective testing in our local environment.

* Corresponding author at: School of Medical Sciences, Bangor University, Bangor, United Kingdom.

E-mail address: Christian.Subbe@Wales.NHS.UK (C.P. Subbe).

2. Methods

Fy Nhaid is a service evaluation that was developed as a collaboration between clinicians from Primary care, Secondary Care and Social Services within the Betsi Cadwaladr University Health board. Fy Nhaid aims to use existing data sources to develop decision rules that help service managers and clinicians to identify patients likely to require support after discharge from hospital.

2.1. Data sources

Electronic point of care (EPOC) is a data-base in the Acute Medical Unit at Wrexham Maelor Hospital that is being used to triage patients on admission to hospital. The data base contains vital signs on admission to hospital, previous medical history in ICD codes, medication as well as a measure of frailty (Clinical Frailty Scale (5)) from over 10,000 patients admitted to the Maelor Hospital. Value of the Clinical Frailty Scale of 5 or more code for patients who are frail.

The post-discharge support data was provided by social services in North-East Wales as part of a data sharing agreement. Data from patients starting new social care packages between 1st April 2014 and 15 February 2016 were included. Social services data comprised the frequency of service use and start and finish data of support packages. Support packages included adaptation of a patient's home, provision of equipment, domiciliary support, nursing care, professional support and supported living and council residential care.

2.2. Principles of evaluation

The EPOC data were housed on a secure server at Betsi Cadwaladr University Health Board (BCUHB). Files were merged with data from social services and anonymized after merger. Anonymized files received a new study specific number. Anonymized data was then transferred to the North Wales Organisation for Randomised Trials in Health, School of Healthcare Sciences (NORTH) at Bangor University, where they were stored on a secure server.

NORTH provides research support to BCUHB. NORTH supported the merger of the files described above as part of the Health and Care Research Wales Clinical Research Centre and Research Infra-structure and Technical Support Group and undertook the primary analysis.

2.3. Ethics

The analysis of data was undertaken as part of a service development program between the department for General, Specialist and Community Medicine that comprises adult inpatient services and primary care provision throughout North Wales and Wrexham Social Services. The process was approved by the local data governance officer.

2.4. Selection of patients

We included patients admitted through the Acute Medical Unit (AMU) at Wrexham Maelor Hospital and seen by social services in the Wrexham area within 60 days after discharge. These were compared to a sample of patients who were admitted to the same unit but were not registered as having received social service support. Controls were matched by post-code.

2.5. Data analysis

Logistic regression analysis with forward and backward selection was used to identify significant predictors of support by social services. Subsequently decision tree learning was applied. The analysis was run using the chi-square automatic interaction detector (CHAID) method [6]. A significance level of 0.05 was applied for

either splitting or merging categories. Additionally cross-validation was conducted.

All analysis was conducted using SPSS version 22.

2.6. Selection of candidate variables

The binary logistic regression analysis included the following variables. With the exception of the number of medications on discharge all data were from the admissions dataset:

- Age in years
- Nursing home admission
- Prior illness
- Diagnosis of diabetes
- Smoking status
- Alcohol excess
- Respiratory rate
- Clinical Frailty Scale [5]
- Systolic blood pressure
- Diastolic blood pressure
- Heart rate
- Simple Clinical Score [7]
- Number of medications as recorded on discharge
- National Early Warning Score (NEWS) [8].

3. Results

The group of patients that required social services comprised 264 patients with a mean age of 80.4 (SD = 9.9) years. The matched group of patients receiving no social services comprised 152 patients with a mean age of 67.5 (SD = 14.2) years.

The variables remaining in the binary logistic regression model following forward and backward selection are listed in Table 1. The forward model achieved a 93.9% classification accuracy and the backward model achieved a 94.1% classification accuracy, based on using the entire dataset. The result demonstrates that important variables are age, Clinical Frailty Scale and visit number. Because NEWS is more widely used this was selected for the classification analysis rather than the Simple Clinical Score.

The decision tree classification analysis was run with all continuous variables left as continuous. The results of this analysis suggested categorizing these variables in the manner shown in Fig. 1. The odds ratios are listed in Table 2. The visit number and the NEWS score have low odds ratios whereas Clinical Frailty Scale and age have higher odds ratios.

Running the decision tree classification resulted in the model shown in Fig. 1. The classification of the model is recorded in Table 3. The model achieved an 85.4% classification accuracy for the training data and 84.1% accuracy for the test data.

Results from the decision rules can be summarized as follows: Social service input after discharge is common in patients with a NEWS value of 0 and age over 80 or significant frailty (CFS > 4) and only one admission.

Table 1

The variables remaining in the binary logistic regression model following forward and backward selection.

Forward model	Backward model
Age	Age
Clinical Frailty Scale	Clinical Frailty Scale
Simple Clinical Score	Heart rate
Visit number (number of admissions)	Visit number (number of admissions)
Constant	NEWS score
	Constant

Download English Version:

<https://daneshyari.com/en/article/8758252>

Download Persian Version:

<https://daneshyari.com/article/8758252>

[Daneshyari.com](https://daneshyari.com)