

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Journal of Acute Disease

journal homepage: www.jadweb.orgOriginal article <http://dx.doi.org/10.1016/j.joad.2016.07.002>

Clinical study of emergency treatment and selective closed reduction for the treatment of supracondylar humerus fracture in children

Q1

Q6 Wei Zhong¹, Xue-Wen Wang^{2*}Q2 ¹Department of Sport Medicine and Orthopedic, Affiliated Hospital of Weifang Medical University, Weifang 261031, Shandong, ChinaQ2 ²Department of Spine, Affiliated Hospital of Weifang Medical University, Weifang 261031, Shandong, China

Q3 ARTICLE INFO

Article history:

Received 14 Jun 2016

Accepted 25 Jul 2016

Available online xxx

Keywords:

Emergency treatment

Selective closed reduction

Supracondylar fracture

Percutaneous fixation

ABSTRACT

Objective: To study the effect of emergency treatment, selective closed reduction combined with percutaneous Kirschner wire fixation on the treatment of Gartland type-II and type-III supracondylar humerus fracture.

Methods: Children who sustained the Gartland type-II and type-III supracondylar fractures of humerus treated with selective closed reduction combined with percutaneous Kirschner wire fixation in our hospital from May 2012 to August 2015 were analyzed retrospectively. They were divided into group A (emergency operation group) and group B (selective operation group) according to different operation timing. Perioperative situation, blood biochemical parameters, swelling degree and elbow joint function of affected limb were compared between two groups.

Results: Operation time for patients of group A was significantly shorter than that of group B [(17.19 ± 2.85) vs. (21.43 ± 3.91) min], and frequency of fluoroscopy during operation of group A was obviously less than that of group B [(6.03 ± 0.95) vs. (7.61 ± 0.92) times]. Swelling index of affected limb in group A at 3 days, 5 days and 7 days after injury was all significantly lower than that in group B [(1.20 ± 0.17) vs. (1.38 ± 0.14), (1.13 ± 0.13) vs. (1.30 ± 0.18), (1.02 ± 0.15) vs. (1.22 ± 0.15)]. Hospital for special surgery score at 1 week, 2 weeks, 3 and 4 weeks after removing Kirschner wire had no significant difference between group A and B (88.75 ± 10.18) vs. (89.14 ± 10.52), (94.22 ± 10.85) vs. (93.85 ± 11.08), (95.52 ± 11.27) vs. (95.92 ± 12.19), (95.43 ± 10.96) vs. (96.02 ± 11.38). Contents of serum alanine transaminase, aspartate aminotransferase, total protein, albumin and C-reactive protein in perioperative period had no obvious difference between patients in group A and B.

Conclusions: Emergency closed reduction combined with percutaneous Kirschner wire fixation for Gartland type-II and type-III supracondylar humerus fracture in children has less trauma, low swelling degree of affected limb in perioperative period, and good effect on the recovery of the elbow joint function after operation.

Q4

1. Introduction

Supracondylar humerus fracture is one of the most common types of fracture in children, of which the Gartland type-I closed

fracture, no obvious displacement with intact anterior and posterior periosteum, can be treated well by manual reduction and plaster external fixation. While the Gartland type-II and type-III closed fractures have a significant displacement leading to a very difficult and complicated clinical procedure^[1,2]. At present, the common clinical therapies for Gartland type-II and type-III fractures include manual reduction, external plaster fixation, open reduction and internal fixation and closed reduction combined with internal Kirschner wire fixation, and so on. The recurrence rate of fracture displacement after operation was high due to the small plane contact areas of broken ends from Gartland type-II and type-III supracondylar humerus fracture, the swelling of soft tissue subsides after fractures reduction, and incoordination of children of hyperactivity in therapy of immobilization^[3,4].

*Corresponding author: Xue-Wen Wang, Department of Sport Medicine and Orthopedic, Affiliated Hospital of Weifang Medical University, Weifang 261031, Shandong, China.

Tel: +86 15954409479

E-mail: 15954409479@163.com

The study protocol was performed according to the Helsinki declaration and approved by the ethic committee of affiliated hospital of Weifang medical university. Informed written consent was obtained from affiliated hospital of Weifang medical university.

Peer review under responsibility of Hainan Medical College. The journal implements double-blind peer review practiced by specially invited international editorial board members.

In recent years, clinical academics have basically reached a consensus on the operation method for Gartland type-II and type-III supracondylar humerus fracture. Closed reduction and percutaneous Kirschner wire fixation has been considered as the preferred treatment, which avoid a great trauma caused by open reduction and internal fixation and also the displacement of fracture ends caused by external plaster fixation^[5,6]. However, the operation timing of closed reduction and percutaneous Kirschner wire fixation for Gartland type-II and type-III supracondylar humerus fracture in children has not yet reached an agreement. In the following study, we compared the effect between emergency treatment and selective closed reduction combined with percutaneous Kirschner wire fixation for Gartland type-II and type-III supracondylar humerus fracture.

2. Materials and methods

2.1. Subjects

Children who sustained the Gartland type-II and type-III supracondylar humerus fracture treated with selective closed reduction combined with percutaneous Kirschner wire fixation in our hospital from May 2012 to August 2015 were analyzed retrospectively, and the inclusion criteria were as follow: (1) their age was from 3 to 12 years; (2) their fracture time was not longer than 8 h; (3) they were suffering from unilateral supracondylar fractures of humerus and diagnosed as Gartland type-II or type-III fractures using imaging test after admission to hospital; (4) they had reached the operative indication and received treatment of closed reduction combined with percutaneous Kirschner wire fixation after admission to hospital; (5) their medical records were complete. Exclusion criteria were as follows: (1) children combined with brain, chest and abdomen injuries; (2) children combined with another parts of fractures; (3) children suffering from open fracture; (4) children with bad terminal blood circulation. A total of 103 cases were recruited and divided into group A (emergency operation group) and group B (selective operation group) according to the medical history review and different operation timing.

2.2. Operation methods

Children of both group A and B were treated by giving vein combined with brachial plexus anesthesia by doctors of the same group. Children from group A took operation after 8 h from injury, while children from group B took operation after 3–5 days from injury. The operation methods were as follows: patients were put down in supine position and the affected limbs were kept a little outstretched; elbow joint was half folded for traction after disinfecting the drapes; epitrochlea was touched with fingers after manual reduction to make sure that the bones were flat and complete without a sense of steps; two Kirschner wires were driven percutaneously into the opposite cortical bone started from lateral condyle until there was a sense of unimpeded. Posterior, anterior and lateral X-ray examination of distal humerus were carried out during operation to make sure that reduction of distal humerus and the placement of Kirschner wire were satisfactory, and then the third Kirschner wire was placed in; X-ray examination was used again to adjust the position of Kirschner wire. The elbow joint flexed to 90° and fixed externally with plaster after operation; the plaster was removed 3

weeks later and the Kirschner wires were removed 4–5 weeks later based on the fracture healing.

2.3. Evaluation of preoperative period situation

Operation time and X-ray examination times of two groups were recorded during operation. The limb swelling index was assessed after 1 day, 3 days, 5 days and 7 days from injury according to the following methods: the circumference of the most obvious swelling part of the affected limbs was measured, and the circumference of same part of opposite unaffected limbs was measured then. The circumference of affected/unaffected limbs was used as the limb swelling index. The samples of peripheral blood were collected and the blood serum was separated at 1 day before and after the operation respectively. Fully automatic biochemical analyzer was used to detect the alanine transaminase (ALT), aspartate aminotransferase (AST), total protein (TP), albumin (Alb) and C-reactive protein (CRP). Hospital for special surgery scoring system (HSS) was used to evaluate the elbow joint function of children in two groups after 1 week, 2 weeks, 3 weeks and 4 weeks from removing the Kirschner wire after operation.

2.4. Statistics methods

Software SPSS version 19.0 was used to analyze the data. Measurement data were expressed as mean \pm SD and analyzed using *t*-test. Enumeration data were described using frequency and analyzed using *Chi*-square. Difference was considered statistically significant when *P* is less than 0.05.

3. Results

3.1. General clinical data of children in two groups

There were 56 cases in emergency operation group (group A) which included 32 cases of boys and 24 cases of girls whose age was [(6.61 \pm 0.92) years]. And there were 35 cases of children diagnosed with Gartland type-III fracture and 21 cases diagnosed with Gartland type-II fracture. The timing from injury to admission to hospital was [(5.86 \pm 0.82) h] and the timing from injury to operation was [(4.94 \pm 0.77) h]. There were 47 cases in selective operation group (group B) which included 28 cases of boys and 19 cases of girls whose ages were [(6.48 \pm 0.83) years]. And there were 30 cases diagnosed with Gartland type-II fracture and 17 cases diagnosed with Gartland type-III fracture. The timing from injury to admission to hospital was [(6.02 \pm 0.79) h] and the timing from injury to operation was [(92.95 \pm 11.38) h]. Both two groups were of no significant difference in gender, age, fracture type and timing from injury to admission to hospital, but the timing from injury to operation of group A was obviously shorter than that of group B (*P* < 0.05) (Table 1).

3.2. Perioperative situation

The operation time from group A was significantly shorter than that of group B [(17.19 \pm 2.85) vs. (21.43 \pm 3.91) min], and frequency of X-ray examination during operation from group A was obviously less than that of group B [(6.03 \pm 0.95) vs. (7.61 \pm 0.92) times] (*P* < 0.05). There was no obvious difference of swelling limbs index between two groups after 1 day from

Download English Version:

<https://daneshyari.com/en/article/8758795>

Download Persian Version:

<https://daneshyari.com/article/8758795>

[Daneshyari.com](https://daneshyari.com)