ARTICLE IN PRESS

NEFROLOGIA 2016; xxx(xx): xxx-xxx







Letter to the Editor

Fibronectin glomerulopathy in a 88 year-old male with acute kidney injury on chronic kidney disease: A case report and a review of the literature

Glomerulopatía fibronectina en un anciano de 88 años de edad con lesión renal aguda en enfermedad renal crónica: reporte de un caso y revisión sistemática de la literatura

Dear Editor,

Fibronectin glomerulopathy (FGP) was first recognized through direct staining in 1992 by Mazzucco et al.¹ It was regarded as a rare autosomal dominant disease that tends to happen in various age groups and affects both sexes. The clinical features of this disease were proteinuria, typically nephrotic range, microscopic hematuria, hypertension and a slow decline in renal function.^{1–11} However, some patients progress to end-stage renal disease (ESRD) and dialysis has been reported.^{5,12–14} Herein we reported a case of FGP that was diagnosed based on the classic ultrastructural features described above. This case was unique in that FGP presents in a male who was 88 years of age and has no known family history of FGP.

An 88-year-old gentleman was admitted with anuria requiring dialysis for uremic encephalopathy and volume overload. He had an extensive past medical history including chronic renal failure (baseline Cr 200–240 μ mol/L) one month ago, hypertension, obstructive sleep apnea, mild cognitive impairment, query previous congestive heart failure, diastolic dysfunction, COPD, macular degeneration, benign prostatic hypertrophy and an anemia that was being investigated as an outpatient. Baseline level of proteinuria was unknown. There was no family history of kidney disease.

Relevant initial blood work included a serum creatinine of $1148\,\mu\text{mol/L}$, urea $82.2\,\text{mmol/L}$ and potassium $5.7\,\text{mmol/L}$. During his stay in hospital, a 24-h urine collection was done and only $100\,\text{mL}$ of urine were collected with $100\,\text{mg}$ of protein and a protein excretion of $1\,\text{g/L}$. The other results of blood examination are as follows: a normal ESR, C3 and C4 values, negative ANA, c-ANCA, p-ANCA, anti-GBM antibody, hepatitis B and C serology, renal ultrasound showed normal size

of the kidney. The extranuclear antigen screen was negative except RNP. Immunofixation revealed a serum IgG M-protein with kappa light chain specificity. Bone marrow studies were unremarkable.

A renal biopsy was performed and light microscopy showed a lobular appearance with cellular mesangial nodules that were expanded by matrix. There was also prominent arteriolar fibrointimal hyperplasia and arteriolar hyalinosis. Congo red staining was negative for amyloid. Immunostaining was 2+ for C3 in a granular pattern in the glomerular capillary walls, however staining was negative for IgA, IgG, IgM, kappa, lambda, albumin, fibrin and C1. Electron microscopy showed prominent expansion of the mesangial and subendothelial regions with numerous randomly arranged fibrils seen under high power lens, measuring 10.7–16.8 nm in diameter. Polyclonal rabbit antibody (DAKO, A0245) was used in staining of fibronectin and there was positivity in the mesangium, mesangial nodules and capillary walls (Fig. 1).

The patient received regular hemodialysis and other supportive treatment including anti-hypertension, improving anemia, and preventing renal osteopathy. Unfortunately, this patient died of heart attack half a year later.

To our knowledge, most cases of FGP have presented and been diagnosed in patients under 65-year-old. As this patient was diagnosed with FGP at the age of 88, we reviewed the cases reported in literatures and attempted to explain how this patient's disease course fits within our understanding of FGP.

A PubMed search using the terms "fibronectin glomerulopathy" yielded 16 articles reports in English of patients with biopsy and fibronectin staining proven FGP. This allowed us to compare the disease features of 64 patients, including the patient described in this paper (Table 1). These patients

Last renal

outcome

stable

stable

N/A

0.82

N/A

stable

1.2

0.8

1.54

3.54

N/A

Dialysis

1 patients:

transplant

transplant 4 patients: dead 1 patient: normal Cr, renal cell carcinoma

2 patients:

hemodalysis

2 patients: CAPD

3 patients: dead

Cr was stable

4 patients: Cr was

2 patients: dialysis

1 patients: dead 1 patient: normal Cr, renal cell carcinoma

Cr was stable

1 patient: Cr was

9 patients: Cr was

4 patients: dialysis 4 patients:

Please cite this article in press as: Cheng G, et al. Fibronectin glomerulopathy in a 88 year-old male with acute kidney injury on chronic kidney disease: A case report and a review of the literature. Nefrologia. 2016. http://dx.doi.org/10.1016/j.nefro.2016.09.015

Note: If a particular piece of information was not available, the space is filled with "N/A".

Download English Version:

https://daneshyari.com/en/article/8774871

Download Persian Version:

https://daneshyari.com/article/8774871

<u>Daneshyari.com</u>