

Accepted Manuscript

Title: Urinary-Cutaneous Fistulae Following Conservative Management of Extraperitoneal Bladder Ruptures

Author: Niels V Johnsen, Rachel Sosland, Melissa R Kaufman, Oscar D Guillamondegui, Roger R Dmochowski

PII: S0090-4295(17)30703-3
DOI: <http://dx.doi.org/doi: 10.1016/j.urology.2017.06.034>
Reference: URL 20532

To appear in: *Urology*

Received date: 31-5-2017
Accepted date: 26-6-2017

Please cite this article as: Niels V Johnsen, Rachel Sosland, Melissa R Kaufman, Oscar D Guillamondegui, Roger R Dmochowski, Urinary-Cutaneous Fistulae Following Conservative Management of Extraperitoneal Bladder Ruptures, *Urology* (2017), <http://dx.doi.org/doi: 10.1016/j.urology.2017.06.034>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Urinary-Cutaneous Fistulae Following Conservative Management of
Extraperitoneal Bladder Ruptures

Running head: Urinary-Cutaneous Fistulae after Bladder Rupture

Niels V Johnsen, Rachel Sosland, Melissa R Kaufman, Oscar D Guillamondegui,
Roger R Dmochowski

Niels Vass Johnsen, MD
Department of Urologic Surgery
A-1302 Medical Center North
Vanderbilt University Medical Center
Nashville, TN 37232
niels.v.johnsen@vanderbilt.edu
njohnsen@gmail.com
FAX: (615) 322-8990
Phone: (504) 250-6694

Word count: 2999
Abstract: 233

Funding: No funding was obtained for completion of this study.

Conflicts of interest: No study author reports any potential conflicts of interest involved with completion of this study.

Abstract

Objective:

To evaluate our experience with urinary-cutaneous fistulae (UCF) in patients following conservative management of extraperitoneal bladder ruptures (EBR) due to blunt trauma.

Materials and Methods:

Patients with blunt-trauma related EBR who underwent primary management with urethral catheter drainage alone from 2000 to 2015 were identified in our institutional trauma registry. The demographics, hospital courses and outcomes

Download English Version:

<https://daneshyari.com/en/article/8776435>

Download Persian Version:

<https://daneshyari.com/article/8776435>

[Daneshyari.com](https://daneshyari.com)