

Accepted Manuscript

Title: Does a human chorionic gonadotropin level of over 20,000 IU/L four weeks after uterine evacuation for complete hydatidiform mole constitute an indication for chemotherapy for gestational trophoblastic neoplasia?

Authors: Antonio Braga, Andressa Biscaro, Jessye Melgarejo do Amaral Giordani, Maurício Viggiano, Kevin M. Elias, Ross S. Berkowitz, Michael J. Seckl

PII: S0301-2115(18)30049-6
DOI: <https://doi.org/10.1016/j.ejogrb.2018.02.001>
Reference: EURO 10220

To appear in: *EURO*

Received date: 10-10-2017
Revised date: 22-1-2018
Accepted date: 1-2-2018

Please cite this article as: Braga Antonio, Biscaro Andressa, do Amaral Giordani Jessye Melgarejo, Viggiano Maurício, Elias Kevin M, Berkowitz Ross S, Seckl Michael J. Does a human chorionic gonadotropin level of over 20,000 IU/L four weeks after uterine evacuation for complete hydatidiform mole constitute an indication for chemotherapy for gestational trophoblastic neoplasia?. *European Journal of Obstetrics and Gynecology and Reproductive Biology* <https://doi.org/10.1016/j.ejogrb.2018.02.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Does a human chorionic gonadotropin level of over 20,000 IU/L four weeks after uterine evacuation for complete hydatidiform mole constitute an indication for chemotherapy for gestational trophoblastic neoplasia?

Antonio Braga MD, PhD ^{1,2*}, Andressa Biscaro MD ²,

Jessye Melgarejo do Amaral Giordani PhD ³, Maurício Viggiano MD ⁴, Kevin M. Elias MD ⁵,

Ross S. Berkowitz MD ⁵, Michael J. Seckl MD, PhD, FRCP ⁶

¹- Rio de Janeiro Trophoblastic Disease Center, Maternity School, Federal University of Rio de Janeiro and Antonio Pedro University Hospital at Fluminense Federal University, Rio de Janeiro, Brazil. Postgraduate Program in Perinatal Health, Maternity School of Rio de Janeiro Federal University.

²- Postgraduate Program in Maternal and Child Health, Fluminense Federal University, Niterói, Brazil.

³- Postgraduate Program in Dentistry, Federal University of Santa Maria, Santa Maria, RS, Brazil.

⁴- Goiás Trophoblastic Disease Center, Goiás Federal University, Goiânia, Brazil.

⁵- New England Trophoblastic Disease Center, Donald P. Goldstein, MD. Trophoblastic Tumor Registry. Department of Obstetrics and Gynecology, Division of Gynecologic Oncology, Brigham and Women's Hospital, Dana Farber Cancer Institute / Harvard Cancer Center, Harvard Medical School. Boston, United States of America.

⁶- Charing Cross Gestational Trophoblastic Disease Centre, Charing Cross Hospital, Imperial College of London, London, UK.

Download English Version:

<https://daneshyari.com/en/article/8778078>

Download Persian Version:

<https://daneshyari.com/article/8778078>

[Daneshyari.com](https://daneshyari.com)