


Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Middle East Fertility Society Journal

journal homepage: www.sciencedirect.com

Original Article

Epidemiology of dysmenorrhea among workers in Upper Egypt; A cross sectional study

Ahmed E. Arafa^a, Yasser Khamis^b, Hanan E. Hassan^c, Nahed M. Saber^c, Ahmed M. Abbas^{d,*}^a Department of Public Health, Faculty of Medicine, Beni-Suef University, Egypt^b Department of Obstetrics and Gynecology, Faculty of Medicine, Beni-Suef University, Egypt^c Department of Maternal and Newborn Health Nursing, Faculty of Nursing, Beni-Suef University, Egypt^d Department of Obstetrics and Gynecology, Faculty of Medicine, Assiut University, Egypt

ARTICLE INFO

Article history:

Received 6 July 2017

Accepted 11 July 2017

Available online xxxxx

Keywords:

Dysmenorrhea
Textile factories
Working women

ABSTRACT

Objective: To investigate the epidemiology of dysmenorrhea amongst women in two textile factories in Beni-Suef, Egypt compared to non-working controls.**Study design:** A cross sectional study.**Setting:** Beni-Suef, Egypt.**Materials and methods:** The study was conducted on 554 working women in two textile factories in Beni-Suef and 1081 non-working women, matched for age and residence. A structured questionnaire was used and a team of data collectors interviewed the women. The questionnaire included questions about the socio-demographic characteristics of women, history of dysmenorrhea during the past 12 months, associating symptoms, pain relief methods, and sources of information.**Results:** The study reported higher rates of dysmenorrhea during the past 12 months (94.6%) amongst the working women, compared to their controls (90.7%) ($p < 0.05$). Backaches, generalized aching and nervousness were the most likely symptoms reported with menstruation 72.2%, 56.3% and 41% respectively. More than half of the working women who experienced dysmenorrhea reported drinking herbal fluids and taking analgesics to alleviate their symptoms, while 43% had to take sick leaves due to their condition. Women in the study group resorted mostly to family members to get information about menstruation and menstrual disturbances.**Conclusions:** Dysmenorrhea is highly prevalent among women working in the textile factories in Beni-Suef. Further research should focus on the adaptive strategies used by women to avoid the negative impacts of dysmenorrhea. Structuring occupational health programs that handle the menstrual disorders amongst the working women, especially in the industrial settings, should be considered.© 2017 Middle East Fertility Society. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Dysmenorrhea, usually referred to as painful menstruation, is defined as a painful, cramping sensation hitting the lower abdomen. It is often accompanied by other symptoms, such as generalized aching, backaches, headache, anorexia, vomiting, nervousness, and breast tenderness, all occurring just before or during the menses [1].

Several risk factors have been linked to dysmenorrhea including hormonal disturbance, exposure to chronic stress, vitamin defi-

ciencies, smoking, unhealthy dietary habits and physical inactivity [2–4].

Previous national and international literatures have investigated the epidemiology of dysmenorrhea in adolescent girls and concluded substantial variations in the incidence of dysmenorrhea [5–11]. However, few studies have explored dysmenorrhea in working women and investigated the relatedness of this condition to industrial work [12–14].

Being a debilitating condition for many women, dysmenorrhea boosts a major negative impact on the females' quality of life, their productivity at work, and healthcare utilization. Dysmenorrhea is associated also with high rates of absenteeism and restriction from regular activities. As a result, dysmenorrhea is responsible for recognizable economic losses attributed to the costs of medications and healthcare, in addition to the decreased productivity [12–15].

Peer review under responsibility of Middle East Fertility Society.

* Corresponding author at: Department of Obstetrics and Gynecology, Assiut University, Women Health Hospital, 71511 Assiut, Egypt.

E-mail address: bmr90@hotmail.com (A.M. Abbas).<http://dx.doi.org/10.1016/j.mefs.2017.07.002>

1110-5690/© 2017 Middle East Fertility Society. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).Please cite this article in press as: A.E. Arafa et al., Epidemiology of dysmenorrhea among workers in Upper Egypt; A cross sectional study, Middle East Fertil Soc J (2017), <http://dx.doi.org/10.1016/j.mefs.2017.07.002>

The physical problems and the financial losses caused by dysmenorrhea are opposed by a limitation in our knowledge regarding the magnitude of the condition among the working females in Egypt, especially those working in industrial settings, which obstacles developing suitable occupational health programs. In this regards, the objective of this study is to investigate the incidence of dysmenorrhea in women working in two textile factories in Beni-Suef, Egypt, compared to non-working women.

2. Materials and methods

In this cross sectional study, all women working in two textile factories in Beni-Suef were invited to participate in the study conducted in the period between December 2016 and March 2017. Only women who were employed for more than one year, younger than 25 years, had no history of pelvic operations, and were not using hormonal contraceptives by the time of the study were allowed to participate. Of the 718 females working in the two textile factories, 554 females were found eligible and participated in the study. Then, we recruited 1081 women, not having work by the time of the study and matching the study group for age and residence, to serve as controls.

The sample size was determined using Epi-Info version 7 Stat Calc, [Center for Disease Control (CDC), WHO], based upon the following criteria; dysmenorrhea rate of 75%, confidence level of 90%, margin of error of 5% and non-response rate of 40%.

All the women who participated in the study, whether in the study group or the control group, were briefed of the purpose of the study with confirming confidentiality of data. The study was approved by the Research Ethics Committee of the Faculty of Medicine, Beni-Suef University.

Data were collected by a trained team of medical students with a supervisor from the Public Health Department using an interview questionnaire. The questionnaire included three sections:

Section I: Included questions about the socio-demographic characteristics, gynaecological age (Calendar age minus menarche age), physical activity, sleeping hours, exposure to smoking, in addition to questions about the length of menstrual cycle and usage of hormonal contraceptives.

Section II: Had questions about whether the women experienced dysmenorrhea during the past 12 months, and the associating clinical symptoms during the same period.

Section III: Questioned about the pain relief methods used by the women who had dysmenorrhea, and if they had to take sick leaves for their complaints. This section included also a question about the main sources of information women resorted to for getting knowledge about menstruation.

The questionnaire was tested for reliability and the Cronbach's alpha was 0.72, while content validity was assessed by a professor of public health and a professor of gynecology and obstetrics.

Data were analyzed using the software, Statistical Package for Social Science (SPSS Inc. Released 2009, PASW Statistics for Windows, version 18.0: SPSS Inc., Chicago, Illinois, USA). Frequency distribution as percentage and descriptive statistics in the form of mean and standard deviation were calculated. Chi-square, *t*-test and correlations were done whenever needed. P values of less than 0.05 were considered significant.

3. Results

The mean age of the women in the study group was 21.6 ± 1.8 (18–25) years and that of controls was 21.6 ± 1.8 (17–25) years, and most of the participants in both groups (74% in study group

and 71.2% in control group) were residing rural areas, with no statistically significant differences between both groups regarding their age or residence (Table 1).

Almost 30% of the women in the study group were married compared to only 5.7% of the control group ($p < 0.05$). Women in the study group were more likely to have illiterate fathers ($p < 0.05$), but the educational level of their mothers showed no differences from their controls ($p > 0.05$) (Table 1).

Compared to their controls, the working women in our study reported recognizably less hours of sleep (7.6 ± 1.4 h/day in the study group versus 8.3 ± 1.7 h/day in the control group) ($p < 0.05$), and more exposure to smoking, however statistically insignificant ($p > 0.05$). Gynaecological age and length of the menstrual cycle did not show considerable differences between both groups ($p > 0.05$) (Table 1).

When the working women and their controls were asked if they had experienced dysmenorrhea throughout the past 12 months, the incidence of dysmenorrhea among women in the study group was statistically higher than their controls, 94.6% and 90.7%, respectively ($p < 0.05$).

Among the working women, 72.2% reported backaches, 56.3% generalized aching, 41% nervousness, 37.9% had acne or flushing, and 30.7% had headaches. Insomnia abdominal distension and dysuria were the least reported complaints by the women in the study group 12.5%, 11.7% and 9.6%, respectively (Table 2).

For women in the study group who experienced dysmenorrhea, drinking herbal fluids and taking analgesics were the most widely used pain relief methods. Heating pads, rest and exercise were tried by few numbers of women. What is really worth pointing out is that 238 (43%) of women who complained of dysmenorrhea had to take sick leaves at least for one day due to their condition (Table 3).

When the working women were questioned about their sources of information about menstrual cycle and disorders, 65.9% reported family members, 24% friends, 14.6% educational institutions, 14.6% TV and social media, and only 6.9% had to ask doctors or nurses.

4. Discussion

Dysmenorrhea comes on the top of the most common gynaecological disorders worldwide, making it a public health concern. In our study, 94.6% of the women working in two textile factories in Beni-Suef reported dysmenorrhea in the past 12 months. Our finding consists with a study conducted on adolescent girls attending nursing schools in Minia and stated a dysmenorrhea rate of 94.4% [10], but relatively higher than results from other national studies; 75% in Mansoura [8] and 66% [16] to 76.1% [9] in Assiut. Reports from international studies showed wide variations in dysmenorrhea rates from only 38.1% in Lebanon [11] and 44.4% in China [17] to 60% in Canada [18], 76% in the USA [19], and 85.1% in Ethiopia [20]. Such major differences could be attributed to many factors. For example, some of the national [8,9] and the international studies [11,19] explored dysmenorrhea among school or university students, making the mean age of their subjects significantly younger than ours. Makhoulf and Abdul Hameed [16], El-Gilany et al. [8], Weissman et al. [19], and Andersch and Milsom [21] reported higher rates of dysmenorrhea in older women.

In addition, women in our study were asked if they experienced dysmenorrhea throughout the past 12 months only, while other studies explored the same condition during different time spans [18–20]. Besides, our study showed that the women in the working group were more likely to be married. In Egypt, married women are often responsible for raising children, managing house issues in addition to many social and cultural burdens which put them in stressful situations most of the time. Exposure to stress is

Download English Version:

<https://daneshyari.com/en/article/8783214>

Download Persian Version:

<https://daneshyari.com/article/8783214>

[Daneshyari.com](https://daneshyari.com)